

ON THE COMPUTATION OF KÄHLER DIFFERENTIALS AND CHARACTERIZATIONS OF GALOIS EXTENSIONS WITH INDEPENDENT DEFECT

STEVEN DALE CUTKOSKY, FRANZ-VIKTOR KUHLMANN AND ANNA RZEPKA

ABSTRACT. For important cases of algebraic extensions of valued fields, we develop presentations of the associated Kähler differentials of the extensions of their valuation rings. We compute their annihilators as well as the associated differentials. We then apply the results to Galois defect extensions of prime degree. Defects can appear in finite extensions of valued fields of positive residue characteristic and are serious obstructions to several problems in positive characteristic. A classification of defects (dependent vs. independent) has been introduced by the second and the third author. It has been shown that perfectoid fields and deeply ramified fields only admit extensions with independent defect. We give several characterizations of independent defect, using ramification ideals, Kähler differentials and traces of the maximal ideals of valuation rings. All of our results are for arbitrary valuations; in particular, we have no restrictions on their rank or value groups.

1. INTRODUCTION

A central goal of this paper is to give for important cases of algebraic extensions of valued fields a presentation of the associated Kähler differentials of the extensions of their valuation rings. This is crucial not only for applications in the present paper, but also in the subsequent paper [6].

By $(L|K, v)$ we denote a field extension $L|K$ where v is a valuation on L and K is endowed with the restriction of v . The valuation ring of v on L will be denoted by \mathcal{O}_L , and that on K by \mathcal{O}_K . Similarly, \mathcal{M}_L and \mathcal{M}_K denote the unique maximal ideals of \mathcal{O}_L and \mathcal{O}_K . The value group of the valued field (L, v) will be denoted by vL , and its residue field by Lv . The value of an element a will be denoted by va , and its residue by av . The **rank** of a valued field (K, v) is the order type of the chain of proper convex subgroups of its value group vK . All of our results are for arbitrary valuations; in particular, we have no restrictions on their rank or value groups.

Date: 29. 9. 2025.

2010 Mathematics Subject Classification. 12J10, 12J25.

Key words and phrases. Kähler differentials, ramification ideal, different, defect extension, Artin-Schreier extension, Kummer extension.

The first author was partially supported by grant DMS 2054394 from NSF of the United States.

The second author was partially supported by Opus grant 2017/25/B/ST1/01815 from the National Science Centre of Poland. He was also supported by a Miller Fellowship during a four weeks visit to the University of Missouri Mathematics Department in 2022; he would like to thank the people at that department for their hospitality.

The authors thank the referee for many helpful corrections and comments. We also thank Sylvy Anscombe, Arno Fehm, Hagen Knaf and Josnei Novacoski for inspiring discussions and helpful feedback.

Ranks higher than 1 appear in a natural way when local uniformization, the local form of resolution of singularities, is studied. Deeply ramified fields of infinite rank appear in model theoretic investigations of the tilting construction, as presented by Jahnke and Kartas in [12]. Therefore, we do not restrict our computations to rank 1, thereby indicating how Kähler differentials and their annihilators, as well as differents, can be computed in higher rank.

By $\Omega_{B|A}$ we denote the Kähler differentials, i.e., the module of relative differentials, when A is a ring and B is an A -algebra. In Section 4.1, we prove:

Theorem 1.1. *Let $L|K$ be an algebraic field extension of degree n and suppose that A is a normal domain with quotient field K and B is a domain with quotient field L such that $A \subset B$ is an integral extension. Suppose that there exist generators $b_\alpha \in B$ of $L|K$, which are indexed by a totally ordered set S , such that $A[b_\alpha] \subset A[b_\beta]$ if $\alpha \leq \beta$ and*

$$\bigcup_{\alpha \in S} A[b_\alpha] = B.$$

Further suppose that there exist $a_\alpha, a_\beta \in A$ such that $a_\beta \mid a_\alpha$ if $\alpha \leq \beta$ and for $\alpha \leq \beta$, there exist $c_{\alpha,\beta} \in A$ and expressions

$$(1) \quad b_\alpha = \frac{a_\alpha}{a_\beta} b_\beta + c_{\alpha,\beta}.$$

Let h_α be the minimal polynomial of b_α over K . Let U and V be the B -ideals

$$U = (a_\alpha \mid \alpha \in S) \text{ and } V = (h'_\alpha(b_\alpha) \mid \alpha \in S).$$

Then we have a B -module isomorphism

$$(2) \quad \Omega_{B|A} \cong U/UV.$$

For the case where $(L|K, v)$ is a valued field extension satisfying condition (8) below and $A = \mathcal{O}_K$ and $B = \mathcal{O}_L$, we compute V in (44) and for arbitrary $\gamma \in S$, we obtain a B -module isomorphism

$$(3) \quad \Omega_{\mathcal{O}_L|\mathcal{O}_K} \cong U/b_\gamma^\dagger U^n \quad \text{with} \quad b_\gamma^\dagger := \frac{h'_\gamma(b_\gamma)}{a_\gamma^{n-1}}.$$

We determine the annihilator of U/UV and thus of $\Omega_{\mathcal{O}_L|\mathcal{O}_K}$ in Proposition 4.7. For information on the different $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K)$, see Theorems 1.6 and 5.4.

We define the defect and defect extensions in Section 2.1. For the construction of a large number of different defect extensions and the role the defect plays in deep open problems in positive characteristic, see [16]. The defect can be understood through jumps in the construction of limit key polynomials and in the construction of pseudo-convergent sequences. A couple of recent references explaining this phenomenon are [25] and [7].

We are generally interested in the study of defect extensions of arbitrary finite degree. As explained in Section 2.1, via ramification theory this can be reduced to the investigation of purely inseparable extensions and of Galois extensions of degree $p = \text{char } Kv > 0$.

We find explicit realizations for the assumptions of Theorem 1.1 for Galois defect extensions of prime degree and deduce the next theorem by combining Theorems 4.2

and 4.3. For the definition of higher ramification groups and ramification ideals, see Section 2.7.

Theorem 1.2. *Let $\mathcal{E} = (L|K, v)$ be a Galois defect extension of prime degree p . If $\text{char } K = 0$, then assume that K contains all p -th roots of unity. Then \mathcal{E} has a unique ramification ideal $I_{\mathcal{E}}$, and there is an \mathcal{O}_L -module isomorphism*

$$(4) \quad \Omega_{\mathcal{O}_L|\mathcal{O}_K} \cong I_{\mathcal{E}}/I_{\mathcal{E}}^p.$$

In [26], Novacoski and Spivakovsky use the theory of key polynomials to derive a presentation of $\Omega_{\mathcal{O}_L|\mathcal{O}_K}$ for finite pure extensions $(L|K, v)$ under the condition $vL = vK$ (which holds for all extensions covered by the above theorem). The presentation of $\Omega_{\mathcal{O}_L|\mathcal{O}_K}$ for Galois defect extensions $(L|K, v)$ is also studied by Thatte in [29, 30].

Galois defect extensions of degree p of valued fields of characteristic $p > 0$ (valued fields of **equal positive characteristic**) have first been classified by the second author in [15]. In [20] the classification was extended to the case of Galois defect extensions of degree p of valued fields of characteristic 0 with residue fields of characteristic $p > 0$ (valued fields of **mixed characteristic**), as follows. Take a Galois defect extension $\mathcal{E} = (L|K, v)$ of prime degree p . For every σ in its Galois group $\text{Gal}(L|K)$, with $\sigma \neq \text{id}$, we set

$$(5) \quad \Sigma_{\sigma} := \left\{ v \left(\frac{\sigma b - b}{b} \right) \mid b \in L^{\times}, \sigma b \neq b \right\}.$$

This set is a final segment of $vL = vK$ and independent of the choice of σ ; we denote it by $\Sigma_{\mathcal{E}}$. The \mathcal{O}_L -ideal $I_{\mathcal{E}} := (a \in L \mid va \in \Sigma_{\mathcal{E}})$ is the unique **ramification ideal** of \mathcal{E} . Detailed information on $\Sigma_{\mathcal{E}}$ and $I_{\mathcal{E}}$ is given in Sections 2.7 and 2.8.

We say that \mathcal{E} has **independent defect** if

$$(6) \quad \left\{ \begin{array}{l} \Sigma_{\mathcal{E}} = \{ \alpha \in vL \mid \alpha > H \} \text{ for some proper convex subgroup } H \\ \text{of } vL \text{ such that } vL/H \text{ has no smallest positive element.} \end{array} \right.$$

If (6) holds, then we will write $H_{\mathcal{E}}$ for H . If there is no such subgroup H , we will say that \mathcal{E} has **dependent defect**. If $vL = vK$ is archimedean (i.e., order isomorphic to a subgroup of \mathbb{R}), then condition (6) just means that $\Sigma_{\mathcal{E}}$ consists of all positive elements in vL and vL has no smallest positive element.

In [11], Gabber and Ramero define deeply ramified fields (K, v) by the property

$$(7) \quad \Omega_{\mathcal{O}_{K^{\text{sep}}|\mathcal{O}_K}} = 0,$$

where K^{sep} denotes the separable-algebraic closure of K . For a valuative definition, which they prove to be equivalent to theirs, and a generalization thereof, see e.g. [20]. In [6] we present a simplified version of their proof. The following is a consequence of the more general part 1) of Theorem 1.10 in [20]:

Theorem 1.3. *Every Galois defect extension of prime degree of a deeply ramified field has independent defect.*

Property (7) together with Theorem 1.3 led us to investigate the connection between Kähler differentials and independent defect. In the present paper we apply

the results stated at the beginning of the introduction to the following situation:

$$(8) \quad \begin{cases} (L|K, v) \text{ is a Galois defect extension of prime degree } p = \text{char } Kv \\ \text{and if } \text{char } K = 0, \text{ then } K \text{ contains all } p\text{-th roots of unity.} \end{cases}$$

In this situation we have $vL = vK$, and $L|K$ is an Artin-Schreier extension if $\text{char } K = p$, and a Kummer extension if $\text{char } K = 0$. We obtain that their defect is independent if and only if $\Omega_{\mathcal{O}_L|\mathcal{O}_K} = 0$; this is the equivalence of assertions a) and d) in Theorem 1.4 below.

Our results for extensions of the form (8) will be complemented in [6] by applying Theorem 1.1 to Galois extensions of prime degree without defect. Using this together with the results of the present paper, it is shown that a valued field is deeply ramified if and only if the Kähler differentials of all Galois extensions of prime degree are zero; moreover, we compute Kähler differentials for all finite Galois extensions satisfying (8).

Another way to characterize independent defect of Galois extensions $(L|K, v)$ of prime degree is through the trace $\text{Tr}_{L|K}(\mathcal{M}_L)$ where $\text{Tr}_{L|K}$ denotes the trace of $L|K$. Note that if $(L|K, v)$ is unbranched, then $\text{Tr}(\mathcal{O}_L) \subseteq \mathcal{O}_K$.

The following theorem summarizes various characterizations of independent defect. The convex subgroups H of vL are in one-to-one correspondence with the coarsenings v_H of v on L in such a way that $v_H L = vL/H$. The maximal ideal of the valuation ring \mathcal{O}_{v_H} of v_H on L is $\mathcal{M}_{v_H} = (a \in L \mid va > H)$. If H is a subgroup of some ordered abelian group Γ , then we call it a **strongly convex subgroup** if it is a proper convex subgroup such that Γ/H has no smallest positive element.

If $\text{char } K = 0$ and $\text{char } Kv = p > 0$, then we denote by $(vK)_{vp}$ the smallest convex subgroup of vK that contains vp (cf. [20]). If $\text{char } K > 0$, then we set $(vK)_{vp} = vK$.

Theorem 1.4. *Assume that the extension $\mathcal{E} = (L|K, v)$ satisfies (8). Then the following assertions are equivalent:*

- a) \mathcal{E} has independent defect,
- b) the ramification ideal $I_{\mathcal{E}}$ of \mathcal{E} is equal to \mathcal{M}_{v_H} for some strongly convex subgroup H of vL ,
- c) $I_{\mathcal{E}}^p = I_{\mathcal{E}}$,
- d) $\Omega_{\mathcal{O}_L|\mathcal{O}_K} = 0$,
- e) $\text{Tr}_{L|K}(\mathcal{M}_L) = \mathcal{M}_{v_H} \cap K$ for some strongly convex subgroup H of vK .

If assertions b) or e) hold, then $H = H_{\mathcal{E}}$, and the valuation ring of v_H is the localization of \mathcal{O}_L with respect to the ramification ideal $I_{\mathcal{E}}$. The value group of the corresponding coarsening of v does not have a smallest positive element.

If vK is archimedean, or more generally, if $(vK)_{vp}$ is archimedean, then $H_{\mathcal{E}}$ can only be equal to $\{0\}$ and if the ramification ideal is a prime ideal, then it can only be equal to \mathcal{M}_L .

If $\Omega_{\mathcal{O}_L|\mathcal{O}_K}$ is annihilated by \mathcal{M}_L , then $\Omega_{\mathcal{O}_L|\mathcal{O}_K} = 0$.

For more equivalent conditions, see part 1) of Proposition 2.10.

The equivalence of assertions a) and b) follows from the definition of independent defect and the fact that $I_{\mathcal{E}} = (a \in L \mid va \in \Sigma_{\mathcal{E}})$. The equivalence of assertions

a) and c) will be proved in part 1) of Proposition 2.10. The equivalence of assertions c) and d) follows from Theorem 1.2. The equivalence of assertions a) and e) follows from the next theorem. The last assertion follows from part 3) of Proposition 4.7. The remaining assertions, in particular the equality $H = H_{\mathcal{E}}$, will be proven together with the next theorem in Section 5.2.

Theorem 1.5. *Assume that the extension $\mathcal{E} = (L|K, v)$ satisfies (8). Then*

$$(9) \quad \mathrm{Tr}_{L|K}(\mathcal{O}_L) = \mathrm{Tr}_{L|K}(\mathcal{M}_L) = (b \in K \mid vb \in (p-1)\Sigma_{\mathcal{E}}) = (I_{\mathcal{E}} \cap K)^{p-1}.$$

The extension \mathcal{E} has independent defect if and only if for some strongly convex subgroup H of vL ,

$$(10) \quad \mathrm{Tr}_{L|K}(\mathcal{O}_L) = \mathrm{Tr}_{L|K}(\mathcal{M}_L) = \mathcal{M}_{v_H} \cap K = (b \in K \mid vb > H).$$

In particular, if $H = \{0\}$ (which is always the case if vK is archimedean, or more generally, if $(vK)_{v_p}$ is archimedean), then this means that

$$\mathrm{Tr}_{L|K}(\mathcal{M}_L) = \mathcal{M}_K.$$

In the mixed characteristic case, if (10) holds, then \mathcal{M}_{v_H} will contain p , so that $\mathrm{char} K v_H = p$.

In [4] the authors introduce the notion of *deeply ramified extensions* $(L|K, v)$ where (K, v) is a local field and $L|K$ is algebraic. It follows from [4, Proposition 2.9] that under the conditions above, $(L|K, v)$ is a deeply ramified extension if and only if $\mathrm{Tr}_{F|L}(\mathcal{M}_F) = \mathcal{M}_L$ for every finite extension $F|L$ (note that as an algebraic extension of a local field, (L, v) is henselian, and so the extension of v to F is uniquely determined); see also [10, Theorem 1.1]. Theorem 1.5 shows that the equality $\mathrm{Tr}_{F|L}(\mathcal{M}_F) = \mathcal{M}_L$ will in general not characterize deeply ramified fields, unless they have rank 1 (as is the case for algebraic extensions of local fields).

The **different** of $(L|K, v)$ is $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) := \mathcal{O}_L :_L \mathcal{C}(\mathcal{O}_L|\mathcal{O}_K)$, where $\mathcal{C}(\mathcal{O}_L|\mathcal{O}_K) := (z \in L \mid \mathrm{Tr}(z\mathcal{O}_L) \subseteq \mathcal{O}_K)$ is the fractional \mathcal{O}_L -ideal called the **complementary ideal**; see [31, Ch. V, §11]. For $b \in \mathcal{O}_L$ and h_b its minimal polynomial over K , the element $h'_b(b) \in \mathcal{O}_L$ is called the **different** of b .

We denote the annihilator of an \mathcal{O}_L -module M by $\mathrm{ann} M$. In Section 5.3, we will prove:

Theorem 1.6. *Assume that the extension $\mathcal{E} = (L|K, v)$ satisfies (8).*

1) *The following \mathcal{O}_L -ideals are equal:*

- i) $I_{\mathcal{E}}^{p-1}$,
- ii) *the ideal V from Theorem 1.1 when applied to $L|K$ with $A = \mathcal{O}_K$ and $B = \mathcal{O}_L$,*
- iii) *the ideal generated by the differentials of all elements of $\mathcal{O}_L \setminus \mathcal{O}_K$,*
- iv) *the ideal generated by the elements of $\mathrm{Tr}_{L|K}(\mathcal{O}_L)$.*

2) *We have that $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = I_{\mathcal{E}}^{p-1}$ if and only if $vI_{\mathcal{E}}^{p-1}$ has no infimum in vL . If $vI_{\mathcal{E}}^{p-1}$ has infimum va in vL for some $a \in L$, then $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = a\mathcal{O}_L \neq I_{\mathcal{E}}^{p-1}$ and $I_{\mathcal{E}}^{p-1} = \mathcal{M}_L \mathcal{D}(\mathcal{O}_L|\mathcal{O}_K)$.*

3) *If (K, v) has rank 1, then $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = \mathrm{ann} \Omega_{\mathcal{O}_L|\mathcal{O}_K}$.*

4) *If the extension \mathcal{E} has independent defect, then $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{O}_L = \mathrm{ann} \Omega_{\mathcal{O}_L|\mathcal{O}_K}$ if $H_{\mathcal{E}} = \{0\}$, and $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{M}_{v_{H_{\mathcal{E}}}} \subsetneq \mathcal{O}_L = \mathrm{ann} \Omega_{\mathcal{O}_L|\mathcal{O}_K}$ otherwise. Conversely,*

if $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K)$ is equal to \mathcal{O}_L or to \mathcal{M}_{v_H} for a nontrivial strongly convex subgroup H of vL , then \mathcal{E} has independent defect.

The reader may note that in rank higher than 1, $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K)$ is not necessarily equal to the annihilator of $\Omega_{\mathcal{O}_L|\mathcal{O}_K}$; this fact has already been observed in [29]. For the computation of annihilators in arbitrary rank, see Section 4.5. For the different of a finite unbranched extension $(L|K, v)$ with $vL = vK$, see Theorem 5.4.

Finally, let us give further characterizations of independent defect, and in particular show that it can be characterized by a simple condition in (K, v) (which is an important point in [12]). If z is an element and S a subset of some valued field (L, v) , then we set

$$v(z - S) := \{v(z - c) \mid c \in S\}.$$

We will write $\wp(X)$ for the Artin-Schreier polynomial $X^p - X$.

Theorem 1.7. *Assume that $\mathcal{E} = (L|K, v)$ is an Artin-Schreier defect extension generated by $\vartheta \in L$ with $\vartheta^p - \vartheta = a \in K$. Then $v(\vartheta - c) < 0$ and $v(a - \wp(c)) = pv(\vartheta - c)$ for all $c \in K$, and therefore,*

$$(11) \quad v(a - \wp(K)) = pv(\vartheta - K).$$

Further, the following are equivalent:

- a) \mathcal{E} has independent defect,
- b) $v(\vartheta - K) = -\{\alpha \in vK \mid \alpha > H\}$ for some strongly convex subgroup H of vK ,
- c) $v(a - \wp(K)) = -\{\alpha \in pvK \mid \alpha > H\}$ for some strongly convex subgroup H of vK ,
- d) there is some strongly convex subgroup H of vK such that for all $b \in K$ such that $vb > H$ there is $c \in K$ satisfying $v(a - \wp(c)) \geq -vb$.

If vK is p -divisible, then these properties are also equivalent to

- e) $pv(\vartheta - K) = v(\vartheta - K)$,
- f) $v(a - \wp(K)) = v(\vartheta - K)$.

If (K, v) has rank 1, then condition d) is equivalent to:

- g) for all $b \in \mathcal{O}_K$ there is $c \in K$ satisfying $v(a - \wp(c)) \geq -vb$.

For the notion of “1-unit”, see Section 2.5.

Theorem 1.8. *Assume that \mathcal{E} is a Kummer defect extension of prime degree p . Then it is generated by a 1-unit $\eta \in L$ with $\eta^p = a \in K$. We have $v(\eta - c) < \frac{1}{p-1}vp$ and $v(a - c^p) = pv(\eta - c)$ for all $c \in K$, and therefore,*

$$(12) \quad v(a - K^p) = pv(\eta - K).$$

Further, the following are equivalent:

- a) \mathcal{E} has independent defect,
- b) $v(\eta - K) = \frac{1}{p-1}vp - \{\alpha \in vK \mid \alpha > H\}$ for some strongly convex subgroup H of vK ,
- c) $v(a - K^p) = \frac{p}{p-1}vp - \{\alpha \in pvK \mid \alpha > H\}$ for some strongly convex subgroup H of vK ,

d) there is some strongly convex subgroup H of vK such that for all $b \in K$ such that $vb > H$ there is $c \in K$ satisfying $v(a - c^p) \geq \frac{p}{p-1}vp - vb$.

If vK is p -divisible, then these properties are also equivalent to

e) $pv(\eta - K) = vp + v(\eta - K)$,

f) $v(a - K^p) = vp + v(\eta - K)$.

If (K, v) has rank 1, then condition d) is equivalent to:

g) for all $b \in \mathcal{O}_K$ there is $c \in K$ satisfying $v(a - c^p) \geq \frac{p}{p-1}vp - vb$.

Remark 1.9. Assume the situation of this theorem. Since K contains a primitive root ζ_p of unity and $v(1 - \zeta_p) = \frac{1}{p-1}vp$, assertion b) is equivalent to

b') $v(\frac{\eta}{1-\zeta_p} - K) = -\{\alpha \in vK \mid \alpha > H\}$ for some strongly convex subgroup H of vK .

We note that conditions g) in the two theorems are elementary in the language of valued rings. It has been pointed out to us by Arno Fehm that in fact also conditions d) are elementary, as it can be stated in the language of valued rings that in the setting of Theorem 1.7, the set

$$\{\alpha \in vK \mid v(a - \wp(K)) < \alpha < -v(a - \wp(K))\}$$

and in the setting of Theorem 1.8, the set

$$\left\{ \alpha \in vK \mid v(a - K^p) - \frac{p}{p-1}vp < \alpha < -v(a - K^p) + \frac{p}{p-1}vp \right\}$$

is a strongly convex subgroup H of vK .

The following can be deduced from Theorems 1.7 and 1.8; for details, see [21]. Take a deeply ramified field (K, v) of rank 1 (e.g. a perfectoid field) with $\text{char } Kv = p > 0$. If it is of equal characteristic, then it satisfies the sentence

$$\forall a, b \in \mathcal{O}_K \exists c \in K : v(a - \wp(c)) \geq -vb ;$$

likewise, if it is of mixed characteristic, then it satisfies the sentence

$$\forall a, b \in \mathcal{O}_K \exists c \in K : v(a - c^p) \geq \frac{p}{p-1}vp - vb .$$

Hence the same holds for every elementary extension (K^*, v^*) of (K, v) . Assume that v^* admits a proper nontrivial coarsening w . Then (K^*, w) satisfies the sentence

$$\forall a \in \mathcal{O}_{(K^*, w)} \exists c \in K^* : w(a - \wp(c)) \geq 0$$

or

$$\forall a \in \mathcal{O}_{(K^*, w)} \exists c \in K^* : w(a - c^p) \geq \frac{p}{p-1}wp ,$$

respectively, which shows that every Artin-Schreier extension or Kummer extension of degree p , respectively, either lies in the henselization of (K^*, w) or is tame (see the definition given in [20]); in particular, they are defectless. This is also a consequence of results obtained in [12], by methods different from those used in [21].

In [21] we will give constructions that will show that all the situations mentioned in the above theorems can appear. To conclude this introduction, let us give an example of an interesting extension with independent defect.

Example 1.10. Choose a prime p and let \mathbb{F}_p denote the field with p elements. Consider the rational function field $\mathbb{F}_p(t)$, equipped with the t -adic valuation v_t , and its perfect hull $K = \mathbb{F}_p(t)(t^{1/p^n} \mid n \in \mathbb{N})$, equipped with the unique extension of v_t . Take a root ϑ of the Artin-Schreier polynomial

$$X^p - X - \frac{1}{t}.$$

The extension $(K(\vartheta)|K, v_t)$ was presented by Shreeram Abhyankar in [1]. It became famous since it shows that there are elements algebraic over $\mathbb{F}_p(t)$ with a power series expansion in which the exponents do not have a common denominator. However, this extension is also an important example for an extension with nontrivial defect. The situation remains the same if we replace $\mathbb{F}_p(t)$ by the field $\mathbb{F}_p((t))$ of formal Laurent series (see [16, Example 3.12]).

In both cases, (K, v_t) is a deeply ramified field, as it is perfect of positive characteristic. Hence by Theorem 1.3, the extension $(K(\vartheta)|K, v_t)$ has independent defect. Now Theorem 1.4 shows that $\Omega_{\mathcal{O}_{K(\vartheta)}|\mathcal{O}_K} = 0$.

2. PRELIMINARIES

2.1. Defect.

A valued field extension $(L|K, v)$ is **unbranched** if the extension of v from K to L is unique. Note that a unbranched extension is automatically algebraic, since every transcendental extension always admits several extensions of the valuation.

If $(L|K, v)$ is a finite unbranched extension, then by the Lemma of Ostrowski [32, Corollary to Theorem 25, Section G, p. 78]),

$$(13) \quad [L : K] = \tilde{p}^\nu \cdot (vL : vK)[Lv : Kv],$$

where ν is a non-negative integer and \tilde{p} the **characteristic exponent** of Kv , that is, $\tilde{p} = \text{char } Kv$ if it is positive and $\tilde{p} = 1$ otherwise. The factor $d(L|K, v) := \tilde{p}^\nu$ is the **defect** of the extension $(L|K, v)$. We call $(L|K, v)$ a **defect extension** if $d(L|K, v) > 1$, and a **defectless extension** if $d(L|K, v) = 1$. Nontrivial defect only appears when $\text{char } Kv = p > 0$, in which case $\tilde{p} = p$. A valued field (K, v) is **henselian** if it satisfies Hensel's Lemma, or equivalently, if all of its algebraic extensions are unbranched. A henselian field (K, v) is called a **defectless field** if all of its finite extensions are defectless.

Throughout this paper, when we talk of a **defect extension** $(L|K, v)$ of **prime degree**, we will always tacitly assume that it is a unbranched extension. Then it follows from (13) that $[L : K] = p = \text{char } Kv$ and that $(vL : vK) = 1 = [Lv : Kv]$; the latter means that $(L|K, v)$ is an **immediate extension**, i.e., the canonical embeddings $vK \hookrightarrow vL$ and $Kv \hookrightarrow Lv$ are onto.

In order to reduce arbitrary finite defect extensions to purely inseparable extensions and Galois extensions of degree $p = \text{char } Kv > 0$, we fix an extension of v from K to its algebraic closure \tilde{K} . The **absolute ramification field of** (K, v) (with respect to the chosen extension of v), denoted by (K^r, v) , is the ramification field of the normal extension $(K^{\text{sep}}|K, v)$. If $a \in \tilde{K}$ is such that $(K(a)|K, v)$ is a defect extension, then $(K^r(a)|K^r, v)$ is a defect extension with the same defect (see [20, Proposition 2.12]). On the other hand, $K^{\text{sep}}|K^r$ is a p -extension (see [15,

Lemma 2.7]), so $K^r(a)|K^r$ is a tower of purely inseparable extensions and Galois extensions of degree p .

2.2. Final segments.

We recall basic notions and facts connected with final segments in ordered abelian groups, and how they relate to elements in valued field extensions. For the details and proofs see Section 2.3 of [15] and Section 3 of [22].

Take a totally ordered set $(T, <)$. For a nonempty subset S of T and an element $t \in T$ we will write $t < S$ if $t < s$ for every $s \in S$, and similarly for $>$ in place of $<$. A set $S \subseteq T$ is called an **initial segment** of T if for each $s \in S$ every $t < s$ also lies in S . Similarly, $S \subseteq T$ is called a **final segment** of T if for each $s \in S$ every $t > s$ also lies in S . By definition, final and initial segments are convex. If S is a final segment of T , then $T \setminus S$ is an initial segment, and vice versa.

For a subset M of T we define

$$M^+ := \{t \in T \mid t > M\} \quad \text{and} \quad M^- := \{t \in T \mid \exists m \in M \ t \geq m\}.$$

Take a subset S in an ordered abelian group Γ , an element $\gamma \in \Gamma$, and $n \in \mathbb{N}_{>0}$. We set $-S := \{-s \mid s \in S\}$, $\gamma + S := \{\gamma + s \mid s \in S\}$, and $nS := \{ns \mid s \in S\}$. We note that if S is a final segment of Γ , then $-S$ is an initial segment of Γ , and $\gamma + S$ is a final segment of Γ . If Γ is divisible by n and S is a final segment of Γ , then also nS is a final segment of Γ . Further, we note that

$$(14) \quad (\gamma + S)^+ = \gamma + S^+ \quad \text{and} \quad (\gamma + S)^- = \gamma + S^-.$$

2.3. Strongly convex subgroups.

Lemma 2.1. *Take an ordered abelian group $(\Gamma, <)$, a nonempty final segment Σ of $\Gamma^{\geq 0} := \{\gamma \in \Gamma \mid \gamma \geq 0\}$, and $m \in \mathbb{N}$, $m \geq 2$. Assume that $\Sigma \neq \Gamma^{\geq 0}$. Then the following assertions are equivalent:*

- a) *there is a strongly convex subgroup Δ of Γ such that $\Sigma = \Gamma^{\geq 0} \setminus \Delta$,*
- b) $\Sigma = (m\Sigma)^-$.

Proof. a) \Rightarrow b): Since Σ is a final segment of $\Gamma^{\geq 0}$, we know that $(m\Sigma)^- \subseteq \Sigma$. Hence in order to prove that $\Sigma = (m\Sigma)^-$, we have to show the reverse inclusion. For this, we will show that for every $\alpha \in \Sigma$ there is $\beta \in \Sigma$ such that $m\beta < \alpha$. The equality $\Sigma = \Gamma^{\geq 0} \setminus \Delta$ implies that $\Sigma = \{\gamma \in \Gamma \mid \gamma > \Delta\}$, hence we have $\alpha + \Delta > 0$ in Γ/Δ . As Γ/Δ has no smallest positive element, the positive elements cannot be bounded away from 0 by an element in the divisible hull of Γ/Δ , and so there must be $\beta \in \Gamma$ such that $0 < \beta + \Delta < \frac{1}{m}(\alpha + \Delta)$. This implies $m\beta + \Delta < \alpha + \Delta$, whence $m\beta < \alpha$, as desired.

b) \Rightarrow a): Since $\Sigma \neq \Gamma^{\geq 0}$ is a nonempty final segment of $\Gamma^{\geq 0}$, we see that $\Gamma^{\geq 0} \setminus \Sigma$ is nonempty and convex. We prove that $\Gamma^{\geq 0} \setminus \Sigma$ is additively closed. Take $\gamma, \gamma' \in \Gamma^{\geq 0} \setminus \Sigma$; we may assume that $\gamma' \leq \gamma$. Hence $0 \leq \gamma + \gamma' \leq m\gamma$. Suppose that $m\gamma \in \Sigma$. Then by the statement of b), there is $\beta \in \Sigma$ such that $m\beta \leq m\gamma$, whence $\beta \leq \gamma$. From the convexity of Σ it follows that $\gamma \in \Sigma$, contradiction. Thus $m\gamma \in \Gamma^{\geq 0} \setminus \Sigma$, and by the convexity of $\Gamma^{\geq 0} \setminus \Sigma$, we obtain that $\gamma + \gamma' \in \Gamma^{\geq 0} \setminus \Sigma$. This proves our claim.

Since $\Gamma^{\geq 0} \setminus \Sigma$ is convex, also $\Delta := \{\gamma, -\gamma \mid \gamma \in \Gamma^{\geq 0} \setminus \Sigma\}$ is convex. Since $\Gamma^{\geq 0} \setminus \Sigma$ is additively closed, it follows by convexity that Δ is additively closed, hence a convex subgroup of Γ .

Take any $\alpha \in \Gamma$ such that $0 < \alpha + \Delta$. It follows that $0 < \alpha \notin \Delta$, hence $\alpha \in \Sigma$. Take $\beta \in \Sigma$ such that $m\beta \leq \alpha$. We have that $\beta + \Delta < m\beta + \Delta$ since otherwise, $(m-1)\beta \in \Delta$ and by convexity, $\beta \in \Delta$, contradiction. Thus $0 < \beta + \Delta < m\beta + \Delta \leq \alpha + \Delta$. This shows that $\alpha + \Delta$ is not the smallest positive element of Γ/Δ . As $\alpha + \Delta$ was arbitrary, we conclude that Γ/Δ has no smallest positive element. \square

2.4. Immediate and unbranched extensions.

Take a valued field (K, v) . A **henselization** of (K, v) is an algebraic extension of (K, v) which admits a valuation preserving embedding in every other henselian extension of (K, v) . Henselizations always exist and are unique up to valuation preserving isomorphism over K ; therefore we will talk of *the* henselization of (K, v) and denote it by $(K, v)^h = (K^h, v^h)$. The henselization of (K, v) is an immediate separable-algebraic extension.

In what follows, we will need the following result, which is [3, Lemma 2.1]:

Lemma 2.2. *A finite extension $(L|K, v)$ is unbranched if and only if $L|K$ is linearly disjoint from some (equivalently, every) henselization of (K, v) .*

Let us consider an immediate but not necessarily algebraic extension $(K(x)|K, v)$. Then by [15, Theorem 2.19] the set $v(x - K) \subseteq vK$ is a final segment of vK ; in particular, it has no maximal element. If $g \in K[X]$ and there is $\alpha \in v(x - K)$ such that for all $c \in K$ with $v(x - c) \geq \alpha$ the value $vg(c)$ is constant, then we will say that **the value of g is ultimately fixed over K** . We call $(K(x)|K, v)$ **pure in x** if the value of every $g(X) \in K[X]$ of degree smaller than $[K(x) : K]$ is ultimately fixed over K . Note that we set $[K(x) : K] = \infty$ if x is transcendental over K .

Lemma 2.3. *Every unbranched immediate extension $(K(x)|K, v)$ of prime degree is pure in x .*

Proof. By the Lemma of Ostrowski, every such extension must have a defect equal to its degree, which must be equal to its residue characteristic $p > 0$.

In the case of a henselian field (K, v) , the assertion follows from [22, Proposition 6.5]. We have to consider the general case. Since $(K(x)|K, v)$ is unbranched, Lemma 2.2 shows that $K(x)|K$ is linearly disjoint from $K^h|K$. Hence $(K^h(x)|K^h, v)$ is a unbranched extension of degree p , and it is immediate since $K^h(x) = K(x)^h$ and henselizations are immediate extensions. By what we have said above, the extension $(K^h(x)|K^h, v)$ is pure.

Further, it follows from [17, Theorem 2] that there is no element $b \in K^h$ such that $v(x - b) > v(x - K)$. Thus, if there were a polynomial $g \in K[X]$ of degree less than p whose value is not ultimately fixed over K , then its value would also not be ultimately fixed over K^h , contradiction. \square

The next lemma follows from [13, Lemma 8] and [22, Lemma 5.2]. Note that if $(K(x)|K, v)$ is an extension such that $v(x - K)$ has no maximal element, then by the proof of [13, Theorem 1], x is limit of a pseudo Cauchy sequence in (K, v)

without limit in K , or equivalently, by [22, part a) of Lemma 4.1] its approximation type over (K, v) is immediate. We use the Taylor expansion

$$(15) \quad f(X) = \sum_{i=0}^n \partial_i f(c)(X - c)^i$$

where $\partial_i f$ denotes the i -th **Hasse-Schmidt derivative** of f .

Lemma 2.4. *Take an immediate extension $(K(x)|K, v)$ that is pure in x . Let p be the characteristic exponent of Kv . Then for every nonconstant polynomial $f \in K[X]$ of degree smaller than $[K(x) : K]$ there is some $\gamma \in v(x - K)$ such that for all $c \in K$ with $v(x - c) \geq \gamma$ and all i with $1 \leq i \leq \deg f$,*

the values $v\partial_i f(c)$ are fixed, equal to $v\partial_i f(x)$, and

the values $v\partial_i f(c) + i \cdot v(x - c)$ are pairwise distinct.

2.5. Artin-Schreier and Kummer extensions.

Every Galois extension of degree p of a field K of characteristic $p > 0$ is an **Artin-Schreier extension**, that is, generated by an **Artin-Schreier generator** ϑ which is the root of an **Artin-Schreier polynomial** $X^p - X - b$ with $b \in K$. For every $c \in K$, also $\vartheta - c$ is an Artin-Schreier generator as its minimal polynomial is $X^p - X - b + c^p - c$. Every Galois extension of prime degree n of a field K of characteristic different from n which contains all n -th roots of unity is a **Kummer extension**, that is, generated by a **Kummer generator** η which satisfies $\eta^n \in K$. For these facts, see [23, Chapter VI, §6].

A **1-unit** in a valued field (K, v) is an element of the form $u = 1 + b$ with $b \in \mathcal{M}_K$; in other words, u is a unit in \mathcal{O}_K with residue 1. Take a Kummer extension $(L|K, v)$ of degree p of fields of characteristic 0 with any Kummer generator η . Assume that $v\eta \in vK$, so that there is $c_1 \in K$ such that $vc_1 = -v\eta$, whence $vc_1\eta = 0$. Assume further that $c_1\eta v \in Kv$, so that there is $c_2 \in K$ such that $c_2v = (c_1\eta v)^{-1}$. Then $vc_2c_1\eta = 0$ and $c_2c_1\eta v = 1$. Furthermore, $K(c_2c_1\eta) = K(\eta)$ and $(c_2c_1\eta)^p = c_2^p c_1^p \eta^p \in K$. Hence $c_2c_1\eta$ is a Kummer generator of $(L|K, v)$ and a 1-unit. This shows that in particular, if $(L|K, v)$ is an immediate Kummer extension, it always admits a Kummer generator which is a 1-unit.

We note that if η is a 1-unit and if $v(\eta - c) > v\eta = 0$, then also c and c^{-1} are 1-units; conversely, if c is a 1-unit, then $v(\eta - c) > 0$.

We will need the following facts.

Lemma 2.5. *Take a valued field (K, v) , $2 \leq n \in \mathbb{N}$, and a primitive n -th root of unity $\zeta \in K$. Then*

$$(16) \quad \prod_{i=1}^{n-1} (1 - \zeta^i) = n.$$

If in addition n is prime, then

$$(17) \quad v(1 - \zeta) = \frac{vn}{n-1}.$$

Proof. Let $f(X) = X^n - 1 = \prod_{i=1}^n (X - \zeta^i)$. Then $f'(X) = nX^{n-1}$ and $\prod_{i=1}^{n-1} (1 - \zeta^i) = f'(1) = n$. This proves equation (16).

If n is prime, then all powers $\zeta^k \neq 1$ are also primitive n -th roots of unity. We may assume that ζ is chosen among the primitive n -th roots of unity such that $v(1 - \zeta)$ is maximal. Take any $k \in \mathbb{N}$ such that $\zeta^k \neq 1$. Then

$$1 - \zeta^k = (1 - \zeta)(1 + \zeta + \dots + \zeta^{k-1}).$$

Since $v\zeta = 0$, we have $v(1 + \zeta + \dots + \zeta^{k-1}) \geq 0$, whence $v(1 - \zeta^k) \geq v(1 - \zeta)$. By our choice of ζ , this yields $v(1 - \zeta^k) = v(1 - \zeta)$. Consequently, all factors of the product in (16) have equal value. This proves equation (17). \square

2.6. Ideals and final segments.

Take a valued field (L, v) . The function

$$(18) \quad v : I \mapsto vI := \{vb \mid 0 \neq b \in I\}$$

is an order preserving bijection from the set of all proper ideals of \mathcal{O}_L onto the set of all final segments of $vL^{>0}$ (including the final segment \emptyset). The set of these final segments is linearly ordered by inclusion, and the function (18) is order preserving: $J \subseteq I$ holds if and only if $vJ \subseteq vI$ holds. The inverse of the above function is the order preserving function

$$(19) \quad \Sigma \mapsto I_\Sigma := (a \in L \mid va \in \Sigma) = \{a \in L \mid va \in \Sigma\} \cup \{0\}.$$

The following facts from general valuation theory are well known:

Lemma 2.6. *The following statements are equivalent:*

a) I_Σ is a prime ideal of \mathcal{O}_L ,

b) $\Sigma = vL^{\geq 0} \setminus H$ for some convex subgroup H of vL .

If b) holds, then $I_\Sigma = \mathcal{M}_{v_H}$ for the coarsening v_H of v on L whose value group is vL/H .

We note that for each $S \subset vL^{\geq 0}$, we have the following equalities of \mathcal{O}_L -ideals:

$$(20) \quad (a \in L \mid va \in S) = (a \in L \mid va \in S^-) = I_{S^-}.$$

Lemma 2.7. *Take a nonempty final segment Σ of $vL^{\geq 0}$ and $m \in \mathbb{N}_{>0}$.*

1) *We have*

$$I_\Sigma^m = (b \in L \mid vb \in m\Sigma) = (b \in L \mid vb \in (m\Sigma)^-).$$

2) *We have $I_\Sigma = I_\Sigma^m$ if and only if $\Sigma = (m\Sigma)^-$.*

Proof. 1): We have

$$\begin{aligned} I_\Sigma^m &= (a^m \mid a \in I_\Sigma) = (a^m \mid a \in L \text{ and } va \in \Sigma) = (b \in L \mid vb \in m\Sigma) \\ &= (b \in L \mid vb \in (m\Sigma)^-), \end{aligned}$$

where the fourth equality follows from (20), and the third equality is seen as follows. The inclusion \subseteq is obvious. Assume that $b \in L$ with $vb \in m\Sigma$, i.e., there is $\alpha \in \Sigma$ such that $m\alpha = vb$. Choose $a \in L$ such that $va = \alpha$. Then $va^m = vb$, hence $b \in (a^m \mid a \in L \text{ and } va \in \Sigma)$. This proves the inclusion \supseteq .

2): This follows immediately from part 1). \square

2.7. Ramification jumps and ramification ideals.

Take a valued field (K, v) . Assume that $L|K$ is a Galois extension, and let $G = \text{Gal}(L|K)$ denote its Galois group. For proper ideals I of \mathcal{O}_L we consider the (upper series of) **higher ramification groups**

$$(21) \quad G_I := \left\{ \sigma \in G \mid \frac{\sigma b - b}{b} \in I \text{ for all } b \in L^\times \right\}$$

(see [32, §12]). Note that $G_{\mathcal{M}_L}$ is the ramification group of $(L|K, v)$. For every ideal I of \mathcal{O}_L , G_I is a normal subgroup of G ([32, (d) on p. 79]). The function

$$(22) \quad \varphi : I \mapsto G_I$$

preserves \subseteq , that is, if $I \subseteq J$, then $G_I \subseteq G_J$. As \mathcal{O}_L is a valuation ring, the set of its ideals is linearly ordered by inclusion. This shows that also the higher ramification groups are linearly ordered by inclusion. Note that in general, φ will neither be injective nor surjective as a function to the set of normal subgroups of G .

Using the function (19), the higher ramification groups can be represented as

$$G_\Sigma := G_{I_\Sigma} = \left\{ \sigma \in G \mid v \frac{\sigma b - b}{b} \in \Sigma \cup \{\infty\} \text{ for all } b \in L^\times \right\},$$

where Σ runs through all final segments of $vL^{>0}$.

Like the function (22), also the function $\Sigma \mapsto G_\Sigma$ is in general not injective. We call Σ a **ramification jump** if

$$\Sigma' \subsetneq \Sigma \Rightarrow G_{\Sigma'} \subsetneq G_\Sigma$$

for all final segments Σ' of $vL^{>0}$. If Σ is a ramification jump, then I_Σ is called a **ramification ideal**. It follows from the definition that an ideal I of \mathcal{O}_L is a ramification ideal if and only if for some subgroup G' of G , I is the smallest ideal such that $G' = G_I$ (cf. [9, §21]).

In this paper we are particularly interested in the case where $\mathcal{E} = (L|K, v)$ is a Galois extension of prime degree p . Then $G = \text{Gal}(L|K)$ is a cyclic group of order p and thus has only one proper subgroup, namely $\{\text{id}\}$, and this subgroup is equal to G_Σ for $\Sigma = \emptyset$. If in this case G itself is the ramification group of the extension, then there must be a unique ramification jump $\Sigma_{\mathcal{E}}$, and we will call $I_{\mathcal{E}} = I_{\Sigma_{\mathcal{E}}}$ *the* ramification ideal of $(L|K, v)$. As we will show in the next section, ramification jump and ramification ideal carry important information about the extension $(L|K, v)$.

2.8. Galois defect extensions of prime degree.

The following is part of Theorem 3.5 of [20]:

Theorem 2.8. *Take a Galois defect extension $\mathcal{E} = (L|K, v)$ of prime degree with Galois group G . Then G is the ramification group of \mathcal{E} . The set Σ_σ does not depend on the choice of the generator σ of G . Writing $\Sigma_{\mathcal{E}}$ for Σ_σ , we have that the final segment $\Sigma_{\mathcal{E}}$ of $vK^{>0}$ is the unique ramification jump of the extension \mathcal{E} . Further, $I_{\mathcal{E}} = I_{\Sigma_{\mathcal{E}}}$ is the unique ramification ideal of the extension \mathcal{E} .*

If $(L|K, v)$ is an Artin-Schreier defect extension with any Artin-Schreier generator ϑ , then

$$(23) \quad \Sigma_{\mathcal{E}} = -v(\vartheta - K).$$

If K contains a primitive root of unity and $(L|K, v)$ is a Kummer extension with Kummer generator η of value 0, then

$$(24) \quad \Sigma_{\mathcal{E}} = \frac{1}{p-1}vp - v(\eta - K).$$

Corollary 2.9. *Let the assumptions be as in the preceding theorem. Then for every $c \in K$,*

$$v(\vartheta - c) < 0 \quad \text{and} \quad v(\eta - c) < \frac{1}{p-1}vp.$$

Proof. Our assertions follow from (23), (24), and the fact that $\Sigma_{\mathcal{E}} \subseteq vK^{>0}$ by Theorem 2.8. \square

Proposition 2.10. *Assume that the extension $\mathcal{E} = (L|K, v)$ satisfies (8).*

1) *The following assertions are equivalent:*

a) *the extension \mathcal{E} has independent defect,*

b) $\Sigma_{\mathcal{E}} = (p\Sigma_{\mathcal{E}})^{-}$,

c) $I_{\mathcal{E}}^p = I_{\mathcal{E}}$.

d) $(I_{\mathcal{E}} \cap K)^p = I_{\mathcal{E}} \cap K$.

If vK is p -divisible, then these properties are also equivalent to $\Sigma_{\mathcal{E}} = p\Sigma_{\mathcal{E}}$.

2) *If \mathcal{E} has independent defect, then the corresponding convex subgroup $H_{\mathcal{E}}$ of vL does not contain vp .*

Proof. 1): By definition, \mathcal{E} has independent defect if and only if $\Sigma_{\mathcal{E}} = vL^{\geq 0} \setminus H_{\mathcal{E}}$ for some strongly convex subgroup $H_{\mathcal{E}}$ of vL . By Lemma 2.1, this holds if and only if $\Sigma_{\mathcal{E}} = (p\Sigma_{\mathcal{E}})^{-}$. By part 2) of Lemma 2.7, this in turn is equivalent to $I_{\mathcal{E}}^p = I_{\mathcal{E}}$ since $I_{\mathcal{E}} = I_{\Sigma_{\mathcal{E}}}$. It is also equivalent to $(I_{\mathcal{E}} \cap K)^p = I_{\mathcal{E}} \cap K$ because $vL = vK$ and therefore, $I_{\mathcal{E}} \cap K = (a \in K \mid va \in \Sigma_{\mathcal{E}})$.

The last assertion of part 1) holds since if vK is p -divisible, then $p\Sigma_{\mathcal{E}}$ is again a final segment, whence $(p\Sigma_{\mathcal{E}})^{-} = p\Sigma_{\mathcal{E}}$.

2): If $\text{char } K = p$, then $vp = \infty$ and the assertion is trivial. In the mixed characteristic case, where \mathcal{E} is a Kummer extension and admits a Kummer generator η of value 0, we know that $0 = v(\eta - 0) \in v(\eta - K)$, hence also $0 \in -v(\eta - K)$. It follows from (24) that $\frac{1}{p-1}vp \in \Sigma_{\mathcal{E}}$. Therefore, $vp \notin H_{\mathcal{E}}$ since otherwise $\frac{1}{p-1}vp \in H_{\mathcal{E}}$ by convexity; but this is a contradiction. \square

3. GENERATION OF IMMEDIATE UNIBRANCHED EXTENSIONS OF VALUATION RINGS

In this section we will assume that $(L|K, v)$ is an immediate extension, and in various cases determine generators for the valuation ring \mathcal{O}_L as an \mathcal{O}_K -algebra.

3.1. The general case.

In this section we consider an immediate but not necessarily algebraic extension $(K(x)|K, v)$. Then the set $v(x - K)$ has no maximal element. For this and the definition of “pure extension”, see Section 2.4.

For every $\gamma \in vK$ we choose $t_\gamma \in K$ such that $vt_\gamma = -\gamma$. For every $c \in \mathcal{O}_K$ we know that $v(x - c) \in vK$ since the extension is immediate, so we may set $t_c := t_{v(x-c)}$ and

$$x_c := t_c(x - c) \in \mathcal{O}_{K(x)}^\times.$$

We use the Taylor expansion (15).

Lemma 3.1. *Assume that the immediate extension $(K(x)|K, v)$ is pure. Then for every $g(x) \in \mathcal{O}_{K(x)} \cap K[x]$ there is $c \in K$ such that $g(x) \in \mathcal{O}_K[x_c]$. If in addition $K(x)|K$ is algebraic, then*

$$\mathcal{O}_{K(x)} = \bigcup_{c \in K} \mathcal{O}_K[x_c].$$

Proof. From Lemma 2.4 we infer that whenever for $c \in K$ the value $v(x - c)$ is large enough, then the values $v\partial_i g(c)(x - c)^i$, $0 \leq i \leq \deg g$, are pairwise distinct. It follows that

$$vg(x) = \min_i v\partial_i g(c)(x - c)^i = \min_i v\partial_i g(c)t_c^{-i}x_c^i.$$

Since $g(x) \in \mathcal{O}_{K(x)}$ and $vx_c = 0$, we find that $\partial_i g(c)t_c^{-i} \in \mathcal{O}_K$ for all i and therefore, $g(x) \in \mathcal{O}_K[x_c]$.

If $K(x)|K$ is algebraic, then $K(x) = K[x]$ and the last assertion of our lemma follows from what we have already proved. \square

Lemma 3.2. *Take $c_1, c_2 \in K$. If $v(x - c_1) = v(x - c_2)$, then $\mathcal{O}_K[x_{c_1}] = \mathcal{O}_K[x_{c_2}]$. If $v(x - c_1) < v(x - c_2)$, then $\mathcal{O}_K[x_{c_1}] \subseteq \mathcal{O}_K[x_{c_2}]$, and if in addition x_{c_1} is integral over K , then $\mathcal{O}_K[x_{c_1}] \subsetneq \mathcal{O}_K[x_{c_2}]$.*

Proof. Assume that $v(x - c_1) = v(x - c_2)$. Then $t_{c_1} = t_{c_2}$ and

$$v(c_1 - c_2) \geq \min\{v(x - c_1), v(x - c_2)\} = -vt_{c_1},$$

whence $t_{c_1}(c_1 - c_2) \in \mathcal{O}_K$. It follows that

$$\mathcal{O}_K[x_{c_1}] = \mathcal{O}_K[x_{c_1} + t_{c_1}(c_1 - c_2)] = \mathcal{O}_K[t_{c_1}(x - c_2)] = \mathcal{O}_K[t_{c_2}(x - c_2)] = \mathcal{O}_K[x_{c_2}].$$

Now assume that $v(x - c_1) < v(x - c_2)$. Then $v(c_1 - c_2) = \min\{v(x - c_1), v(x - c_2)\} = v(x - c_1) = -vt_{c_1} < -vt_{c_2}$, whence $t_{c_1}/t_{c_2} \in \mathcal{M}_K$ and $t_{c_1}(c_1 - c_2) \in \mathcal{O}_K^\times$. This shows that

$$(25) \quad x_{c_1} = \frac{t_{c_1}}{t_{c_2}} \cdot t_{c_2}(x - c_2) + t_{c_1}(c_2 - c_1) \in \mathcal{O}_K[x_{c_2}],$$

whence $\mathcal{O}_K[x_{c_1}] \subseteq \mathcal{O}_K[x_{c_2}]$.

Suppose that $x_{c_2} \in \mathcal{O}_K[x_{c_1}]$ and x_{c_1} is integral over K . Then x_{c_2} can be written in the form

$$x_{c_2} = a_0 + a_1x_{c_1} + a_2x_{c_1}^2 + \dots + a_{p-1}x_{c_1}^{p-1}$$

with $n = [K(x) : K]$ and coefficients in \mathcal{O}_K . Since the powers of x_{c_1} appearing in this expression form a basis of $K(x)|K$, this representation of x_{c_2} is unique even

if one allows the coefficients to be in K . However, according to (25) we can write $x_{c_1} = ax_{c_2} + b$, where $a \in \mathcal{M}_K$ and $b \in \mathcal{O}_K^\times$. This yields the unique representation

$$x_{c_2} = a^{-1}(x_{c_1} - b),$$

which does not have its coefficients in \mathcal{O}_K . This contradiction shows that $\mathcal{O}_K[x_{c_1}] \neq \mathcal{O}_K[x_{c_2}]$. \square

From Lemmas 3.1 and 3.2, we obtain:

Proposition 3.3. *Assume that the immediate algebraic extension $(K(x)|K, v)$ is pure. Then the rings $\mathcal{O}_K[x_c]$, $c \in \mathcal{O}_K$, form a chain under inclusion whose union is $\mathcal{O}_{K(x)}$.*

3.2. The case of immediate Artin-Schreier and Kummer extensions.

An extension of prime degree is a defect extension if and only if it is immediate and unbranched.

Theorem 3.4. *1) Assume that $(L|K, v)$ is an Artin-Schreier defect extension with Artin-Schreier generator ϑ . The rings $\mathcal{O}_K[\vartheta_c]$, where $\vartheta_c = t_{v(\vartheta-c)}(\vartheta - c)$ and c runs through all elements of K , form a chain under inclusion whose union is \mathcal{O}_L . The same holds for the rings $\mathcal{O}_K[t_{v\vartheta}\vartheta]$ when ϑ runs through all Artin-Schreier generators of the extension.*

For $c_1, c_2 \in K$,

$$(26) \quad \mathcal{O}_K[\vartheta_{c_1}] \subsetneq \mathcal{O}_K[\vartheta_{c_2}] \Leftrightarrow v(\vartheta - c_1) < v(\vartheta - c_2).$$

2) Assume that $(L|K, v)$ is a Kummer defect extension of degree p and that η is a Kummer generator which is a 1-unit. The rings $\mathcal{O}_K[\eta_c]$, where $\eta_c = t_{v(\eta-c)}(\eta - c)$ and c runs through all 1-units of K , form a chain under inclusion whose union is \mathcal{O}_L . If $c_1, c_2 \in K$ are 1-units, then

$$(27) \quad \mathcal{O}_K[\eta_{c_1}] \subsetneq \mathcal{O}_K[\eta_{c_2}] \Leftrightarrow v(\eta - c_1) < v(\eta - c_2).$$

Proof. From Lemma 2.3 and Proposition 3.3 we know that \mathcal{O}_L is the union of the chain $(\mathcal{O}_K[\vartheta_c])_{c \in K}$ in the Artin-Schreier extension case, and of the chain $(\mathcal{O}_K[\eta_c])_{c \in K}$ in the Kummer extension case. Further, (26) and (27) follow from Lemma 3.2.

It remains to prove the additional assertion for the Artin-Schreier extension case. By [15, Lemma 2.26], $\vartheta' \in L$ is another Artin-Schreier generator of $L|K$ if and only if $\vartheta' = i\vartheta - c$ for some $i \in \mathbb{F}_p^\times$ and $c \in K$. In particular, $\vartheta - c$ is an Artin-Schreier generator for all $c \in K$. Moreover, if $\vartheta' = i\vartheta - c$, then $v\vartheta' = v(i\vartheta - c) = v(\vartheta - i^{-1}c)$ and therefore, $t_{v\vartheta'} = t_{i^{-1}c}$ and

$$\mathcal{O}_K[t_{v\vartheta'}\vartheta'] = \mathcal{O}_K[t_{v\vartheta'}i^{-1}\vartheta'] = \mathcal{O}_K[t_{v(\vartheta-i^{-1}c)}(\vartheta - i^{-1}c)] = \mathcal{O}_K[\vartheta_{i^{-1}c}].$$

This shows that the ring $\mathcal{O}_K[t_{v\vartheta'}\vartheta']$ is already a member in the chain constructed above and the union is \mathcal{O}_L . \square

Let us consider the **Artin-Schreier case** a bit further, keeping the assumptions of part 1) of Theorem 3.4. We set $t_c := t_{v(\vartheta-c)}$. Take any elements $c_1, c_2 \in K$. It follows from (25) that

$$(28) \quad \vartheta_{c_1} = \frac{t_{c_1}}{t_{c_2}}\vartheta_{c_2} + t_{c_1}(c_2 - c_1).$$

If $v(\vartheta - c_1) \leq v(\vartheta - c_2)$, then $t_{c_1}/t_{c_2} \in \mathcal{O}_K$ and $t_{c_1}(c_2 - c_1) \in \mathcal{O}_K$, as shown in the proof of Lemma 3.2.

Now we turn to the **Kummer case**, keeping the assumptions of part 2) of Theorem 3.4. We choose a primitive p -th root ζ_p of unity and set

$$\chi := (\zeta_p - 1)^{-1}.$$

According to equation (17), the element χ satisfies:

$$(29) \quad v\chi = -\frac{1}{p-1}vp.$$

Hence by Corollary 2.9,

$$v\chi(\eta - c) < 0$$

for all $c \in K$. In the following we will use the abbreviations

$$T_c := t_{v\chi(\eta-c)} \quad \text{and} \quad \eta_{\langle c \rangle} := T_c\chi(\eta - c)$$

in order to simplify our formulas; we observe that

$$(30) \quad vT_c = \frac{1}{p-1}vp - v(\eta - c) > 0.$$

Corollary 3.5. *Assume that $(L|K, v)$ is a Kummer defect extension of degree p . Then for every Kummer generator η which is a 1-unit, the rings $\mathcal{O}_K[\eta_{\langle c \rangle}]$, where c runs through all 1-units of K , form a chain under inclusion whose union is \mathcal{O}_L .*

We have that

$$(31) \quad \mathcal{O}_K[\eta_{\langle c \rangle}] = \mathcal{O}_K[\eta_c].$$

Hence if c_1 and c_2 are 1-units, then

$$(32) \quad \mathcal{O}_K[\eta_{\langle c_1 \rangle}] \subsetneq \mathcal{O}_K[\eta_{\langle c_2 \rangle}] \Leftrightarrow v(\eta - c_1) < v(\eta - c_2).$$

Proof. It suffices to prove (31). We compute:

$$v \frac{T_c}{\chi^{-1}t_c} = v \frac{t_{v\chi(\eta-c)}}{t_{v\chi}t_{v(\eta-c)}} = 0.$$

Hence,

$$\mathcal{O}_K[\eta_{\langle c \rangle}] = \mathcal{O}_K \left[\frac{T_c}{\chi^{-1}t_c} \chi^{-1}t_c \chi(\eta - c) \right] = \mathcal{O}_K[t_c(\eta - c)] = \mathcal{O}_K[\eta_c].$$

Now the assertion of our corollary follows from part 2) of Theorem 3.4. \square

Take any elements $c_1, c_2 \in K$. Similarly as in the Artin-Schreier case, one derives the equation

$$(33) \quad \eta_{\langle c_1 \rangle} = \frac{T_{c_1}}{T_{c_2}}\eta_{\langle c_2 \rangle} + T_{c_1}\chi(c_2 - c_1).$$

Assume that $v(\eta - c_1) \leq v(\eta - c_2)$. Then $vT_{c_1} \geq vT_{c_2}$ and

$$v(c_2 - c_1) \geq \min\{v(\eta - c_1), v(\eta - c_2)\} = v(\eta - c_1) = -vT_{c_1}\chi,$$

whence $T_{c_1}/T_{c_2} \in \mathcal{O}_K$ and $T_{c_1}\chi(c_2 - c_1) \in \mathcal{O}_K$.

3.3. Values of derivatives of minimal polynomials.

In the case of Artin-Schreier extensions and Kummer extensions $(K(x)|K, v)$ of prime degree we have sufficient information about the minimal polynomials f of the various generators x we have worked with in the previous sections, or equivalently, about their conjugates, to work out the values $vf'(x)$. In order to do this, we can either compute f' explicitly, or we can use the formula

$$(34) \quad f'(x) = \prod_{\sigma \in G \setminus \{\text{id}\}} (x - \sigma x),$$

where G is the Galois group of $K(x)|K$.

We keep the notations from the previous sections.

3.3.1. Artin-Schreier extensions.

Take an Artin-Schreier polynomial f with ϑ as its root. Then $f(X) = X^p - X - \vartheta^p + \vartheta$ and $f'(X) = -1$, whence

$$(35) \quad f'(\vartheta) = -1.$$

This is also obtained from (34) since $\{\vartheta - \sigma\vartheta \mid \sigma \in G \setminus \{\text{id}\}\} = \mathbb{F}_p^\times$ and the product of all elements of \mathbb{F}_p^\times is -1 .

For $t \in K^\times$, denote by f_t the minimal polynomial of $t\vartheta$. Then

$$(36) \quad f'_t(t\vartheta) = \prod_{\sigma \in G \setminus \{\text{id}\}} (t\vartheta - \sigma t\vartheta) = t^{p-1} f'(\vartheta) = -t^{p-1}.$$

Lemma 3.6. *Take an Artin-Schreier defect extension $\mathcal{E} = (L|K, v)$ of prime degree p with Artin-Schreier generator ϑ . Denote by g_c the minimal polynomial of ϑ_c . Then the \mathcal{O}_L -ideal $(g'_c(\vartheta_c) \mid c \in K)$ is equal to*

$$I^{p-1}$$

where

$$(37) \quad I = (t_c \mid c \in K) = (a \in L \mid va \in -v(\vartheta - K)) = I_{\mathcal{E}}$$

is the ramification ideal of the extension $(L|K, v)$.

Proof. Applying (36) to $\vartheta_c = t_c(\vartheta - c)$ and keeping in mind that $\vartheta - c$ is an Artin-Schreier generator, we obtain:

$$(38) \quad g'_c(\vartheta_c) = -t_c^{p-1}.$$

This shows that $(g'_c(\vartheta_c) \mid c \in K) = (t_c \mid c \in K)^{p-1} = I^{p-1}$.

The second equality in (37) holds since $vt_c = -v(\vartheta - c)$ by definition. The third equality follows from equation (23) of Theorem 2.8. \square

3.3.2. *Kummer extensions.*

Take $f(X) = X^p - \eta^p$. Then $f'(X) = pX^{p-1}$, whence

$$(39) \quad f'(\eta) = p\eta^{p-1}.$$

This is also obtained from equations (34) and (16) using that $\{\sigma\eta \mid \sigma \in G\} = \{\zeta_p^i \eta \mid i \in \{0, \dots, p-1\}\}$.

Lemma 3.7. *Take a Kummer defect extension $\mathcal{E} = (L|K, v)$ of prime degree p with Kummer generator η which is a 1-unit. Denote by f_c the minimal polynomial of $\eta_{\langle c \rangle}$ over K . Then the \mathcal{O}_L -ideal $(f'_c(\eta_{\langle c \rangle}) \mid c \in K \text{ a 1-unit})$ is*

$$I^{p-1}$$

where

$$(40) \quad I = (T_c \mid c \in K) = \left(a \in L \mid va \in \frac{1}{p-1}vp - v(\eta - K) \right) = I_{\mathcal{E}}$$

is the ramification ideal of the extension $(L|K, v)$.

Proof. We use (34) to compute:

$$f'_c(\eta_{\langle c \rangle}) = \prod_{\sigma \in G} (\eta_{\langle c \rangle} - \sigma\eta_{\langle c \rangle}) = \prod_{\sigma \in G} T_c \chi(\eta - \sigma\eta) = (T_c \chi \eta)^{p-1} \prod_{i=1}^{p-1} (1 - \zeta_p^i),$$

whence by equations (16) and (29), and the fact that $v\eta = 0$,

$$(41) \quad f'_c(\eta_{\langle c \rangle}) = p(T_c \chi \eta)^{p-1} = T_c^{p-1} u$$

with u a unit in \mathcal{O}_L . This shows that

$$(f'_c(\eta_{\langle c \rangle}) \mid c \in K \text{ a 1-unit}) = (T_c \mid c \in K)^{p-1} = I^{p-1}.$$

The second equality in (40) follows from (30). The assertion that $I = I_{\mathcal{E}}$ in the Kummer case follows from equation (24) of Theorem 2.8. \square

4. KÄHLER DIFFERENTIALS AND THEIR ANNIHILATORS FOR ALGEBRAIC FIELD EXTENSIONS

4.1. Proof of Theorem 1.1.

Let $L|K$ be an algebraic field extension. Let $A \subseteq K$ be a normal domain whose quotient field is K . Assume that $z \in L$ is integral over A and let $f(X)$ be the minimal polynomial of z over K . Then $f(X) \in A[X]$ (see [31, Theorem 4, page 260]). Since $f(X)$ is monic, $(f(X)K[X]) \cap R[X] = f(X)R[X]$, so $A[z] \cong A[X]/(f(X))$. Thus,

$$(42) \quad \Omega_{A[z]|A} \cong [A[X]/(f(X), f'(X))]dX \cong [A[z]/(f'(z))]dX$$

by [24, Example 26.J, page 189] and [24, Theorem 58, page 187]. There is a canonical derivation $d_{A[z]|A} : A[z] \rightarrow \Omega_{A[z]|A}$ defined by $g(z) \mapsto [g'(z)]dX$ for $g(X) \in A[X]$, where $[g'(z)]$ is the class of $g'(z)$ in $A[z]/(f'(z))$.

We will now proceed under the assumptions of Theorem 1.1. Before proving the theorem, we derive a formula that we will need in its proof. Let $n = [L : K]$. Then

for all $\alpha \in S$, $n = \deg h_\alpha(X)$ where $h_\alpha(X)$ is the minimal polynomial of b_α over K . Suppose that $\alpha < \beta$. Then

$$\left(\frac{a_\beta}{a_\alpha}\right)^n h_\alpha\left(\frac{a_\alpha}{a_\beta}X + c_{\alpha,\beta}\right)$$

is a monic polynomial of degree n in $K[X]$ which has b_β as a root, so it is the minimal polynomial $h_\beta(X)$ of b_β . By the chain rule,

$$\frac{dh_\beta}{dX_\beta}(b_\beta) = \left(\frac{a_\beta}{a_\alpha}\right)^n \frac{dh_\alpha}{dX}(b_\alpha) \left(\frac{a_\alpha}{a_\beta}\right),$$

and so

$$(43) \quad h'_\alpha(b_\alpha) = \left(\frac{a_\alpha}{a_\beta}\right)^{n-1} h'_\beta(b_\beta).$$

Proof of Theorem 1.1:

From the natural A -algebra isomorphisms $A[b_\alpha] \cong A[X_\alpha]/(h_\alpha(X_\alpha))$, where $h_\alpha(X_\alpha)$ is the minimal polynomial of b_α over K , we have that for $\alpha \leq \beta$, the natural inclusion of A -algebras $A[b_\alpha] \subset A[b_\beta]$ is induced by sending b_α to $\frac{a_\alpha}{a_\beta}b_\beta + c_{\alpha,\beta}$.

By (42), for $\alpha \in S$ we have a natural isomorphism of $A[b_\alpha]$ -modules

$$\Omega_{A[b_\alpha]|A} \cong [A[b_\alpha]/(h'_\alpha(b_\alpha))] dX_\alpha,$$

and so

$$\Omega_{A[b_\alpha]|A} \otimes_{A[b_\alpha]} B \cong [B/(h'_\alpha(b_\alpha))] dX_\alpha.$$

For $\alpha < \beta$, by the universal property of derivations (Proposition on page 182 [24]), there is a unique B -module homomorphism

$$\lambda_{\alpha,\beta} : \Omega_{A[b_\alpha]|A} \otimes_{A[b_\alpha]} B \rightarrow \Omega_{A[b_\beta]|A} \otimes_{A[b_\beta]} B$$

such that there is a commutative diagram of B -module homomorphisms

$$\begin{array}{ccc} \Omega_{A[b_\alpha]|A} \otimes_{A[b_\alpha]} B & \xrightarrow{\lambda_{\alpha,\beta}} & \Omega_{A[b_\beta]|A} \otimes_{A[b_\beta]} B \\ \uparrow & & \uparrow \\ A[b_\alpha] \otimes_{A[b_\alpha]} B & \rightarrow & A[b_\beta] \otimes_{A[b_\beta]} B \end{array}$$

where the vertical arrows are the respective maps

$$d_{A[b_\alpha]/A} \otimes 1 : A[b_\alpha] \otimes_{A[b_\alpha]} B \rightarrow \Omega_{A[b_\alpha]|A} \otimes_{A[b_\alpha]} B$$

and

$$d_{A[b_\beta]/A} \otimes 1 : A[b_\beta] \otimes_{A[b_\beta]} B \rightarrow \Omega_{A[b_\beta]|A} \otimes_{A[b_\beta]} B.$$

Suppose that $z \in A[b_\alpha]$. Then there exists $g(X_\alpha) \in A[X_\alpha]$ such that $z = g(b_\alpha)$, hence $d_{A[b_\alpha]/A}(z) = \frac{dg}{dX_\alpha}(b_\alpha)dX_\alpha$. We have that $z = \left(g\left(\frac{a_\alpha}{a_\beta}X_\beta + c_{\alpha,\beta}\right)\right)(b_\beta)$ so that by the chain rule,

$$d_{A[b_\beta]/A}(z) = \left(\frac{d}{dX_\beta}g\left(\frac{a_\alpha}{a_\beta}X_\beta + c_{\alpha,\beta}\right)\right)(b_\beta)dX_\beta = \frac{dg}{dX_\alpha}(b_\alpha)\frac{a_\alpha}{a_\beta}dX_\beta.$$

Thus $\lambda_{\alpha,\beta}$ is the B -module homomorphism defined by mapping dX_α to $\frac{a_\alpha}{a_\beta}dX_\beta$.

In order to compute the direct limit of the directed system of B -module homomorphisms $\lambda_{\alpha,\beta} : \Omega_{A[b_\alpha]|A} \otimes_{A[b_\alpha]} B \rightarrow \Omega_{A[b_\beta]|A}$ for $\alpha < \beta$, we will introduce an equivalent directed system, which is a little simpler. For $\alpha \in S$, let M_α be the

B -module $M_\alpha = B/(h'_\alpha(b_\alpha))$. The M_α are a family of B -modules. We have isomorphisms of B -modules $\tau_\alpha : \Omega_{A[b_\alpha]|A} \otimes_{A[b_\alpha]} B \rightarrow M_\alpha$ defined by mapping dX_α to 1.

For all $\alpha, \beta \in S$ with $\alpha < \beta$, we have commutative diagrams of B -module homomorphisms

$$\begin{array}{ccc} \Omega_{A[b_\alpha]|A} \otimes_{A[b_\alpha]} B & \xrightarrow{\lambda_{\alpha,\beta}} & \Omega_{A[b_\beta]|A} \otimes_{A[b_\beta]} B \\ \downarrow \tau_\alpha & & \downarrow \tau_\beta \\ M_\alpha & \xrightarrow{\tau_\beta \lambda_{\alpha,\beta} \tau_\alpha^{-1}} & M_\beta. \end{array}$$

and we see that $\tau_\beta \lambda_{\alpha,\beta} \tau_\alpha^{-1}$ is just multiplication by $\frac{a_\alpha}{a_\beta}$. Thus the directed systems

$$\Omega_{A[b_\alpha]|A} \otimes_{A[b_\alpha]} B \xrightarrow{\lambda_{\alpha,\beta}} \Omega_{A[b_\beta]|A} \otimes_{A[b_\beta]} B \text{ for } \alpha < \beta \quad \text{and} \quad M_\alpha \xrightarrow{\frac{a_\alpha}{a_\beta}} M_\beta \text{ for } \alpha < \beta$$

are equivalent.

We have isomorphisms of B -modules

$$\Omega_{B|A} \cong \lim_{\rightarrow} [\Omega_{A[b_\alpha]|A} \otimes_{A[b_\alpha]} B] \cong \lim_{\rightarrow} M_\alpha$$

by [8, Corollary 16.7, page 394], where the direct limits are over $\alpha \in S$.

Write $M_\alpha = R_\alpha/T_\alpha$ where R_α is the directed system of B -modules $R_\alpha = B$ for $\alpha \in S$, with B -module homomorphisms $\frac{a_\alpha}{a_\beta} : R_\alpha \rightarrow R_\beta$ for $\alpha < \beta$, and T_α is the directed system of B -modules $T_\alpha = h'_\alpha(b_\alpha)R_\alpha$ for $\alpha \in S$, with B -module homomorphisms $\frac{a_\alpha}{a_\beta} : T_\alpha \rightarrow T_\beta$ for $\alpha \leq \beta$. We have short exact sequences of B -modules

$$0 \rightarrow T_\alpha \rightarrow R_\alpha \rightarrow M_\alpha \rightarrow 0$$

which are compatible with multiplication by $\frac{a_\alpha}{a_\beta}$ for $\alpha < \beta$. Thus

$$\lim_{\rightarrow} M_\gamma \cong \lim_{\rightarrow} R_\gamma / \lim_{\rightarrow} T_\gamma$$

by [27, Theorem 2.18].

We now compute $\lim_{\rightarrow} R_\gamma$. For $\alpha \in S$, we have B -module homomorphisms $R_\alpha \xrightarrow{a_\alpha} B$ which give commutative diagrams

$$\begin{array}{ccc} R_\alpha & \xrightarrow{\frac{a_\alpha}{a_\beta}} & R_\beta \\ a_\alpha \searrow & & \swarrow a_\beta \\ & B & \end{array}$$

for $\alpha < \beta$. By the universal property of direct limits ([5, Proposition 11.1] or [2, Exercise 18, page 33]) there exists a unique B -module homomorphism $\Psi : \lim_{\rightarrow} R_\gamma \rightarrow B$ giving commutative diagrams of B -module homomorphisms

$$\begin{array}{ccc} R_\alpha & & \\ \frac{a_\alpha}{a_\beta} \downarrow & \searrow \phi_\alpha & \\ R_\beta & \xrightarrow{\phi_\beta} & \lim_{\rightarrow} R_\gamma \\ a_\beta \downarrow & \swarrow \Psi & \\ B & & \end{array}$$

for $\alpha < \beta$. Here $\phi_\alpha : R_\alpha \rightarrow \lim_{\rightarrow} R_\gamma$ are the canonical maps of the direct product.

As we will now show, it follows from some standard properties of direct limits ([5, Proposition 11.3] or [2, Exercise 15, page 33]) and some diagram chasing that as B -ideals,

$$\lim_{\rightarrow} R_\gamma \cong (a_\alpha \mid \alpha \in S) = U.$$

Suppose that $u \in \lim_{\rightarrow} R_\gamma$. Then there exist $\alpha \in S$ and $b \in R_\alpha = B$ such that $u = \phi_\alpha(b)$. Thus $\Psi(u) = a_\alpha b \in U$. Given $\alpha \in S$, $\Psi(\phi_\alpha(1)) = a_\alpha$. Thus the image of Ψ is U . Suppose that $0 \neq u \in \lim_{\rightarrow} R_\gamma$. Then there exist $\alpha \in S$ and $0 \neq b \in R_\alpha$ such that $\phi_\alpha(b) = u$ so $\Psi(u) = a_\alpha b \neq 0$. Thus Ψ is an isomorphism and so

$$\lim_{\rightarrow} R_\gamma \cong U.$$

Similarly,

$$\lim_{\rightarrow} T_\gamma \cong (a_\alpha h'_\alpha(b_\alpha) \mid \alpha \in S).$$

By (43), we have that for $\alpha, \beta < \gamma$,

$$a_\alpha h'_\beta(b_\beta) = \frac{a_\alpha}{a_\gamma} \left(\frac{a_\beta}{a_\gamma} \right)^{n-1} a_\gamma h'_\gamma(b_\gamma).$$

Thus $(a_\alpha h'_\alpha(b_\alpha) \mid \alpha \in S) = UV$ and so $\Omega_{B|A} \cong U/UV$. \square

If we choose any $\gamma \in S$, then we will still have

$$\bigcup_{\gamma \leq \alpha \in S} A[b_\alpha] = B$$

since $(A[b_\alpha])_{\alpha \in S}$ is an increasing chain. Hence we can always assume that S has a minimal element γ .

From now on we consider the case of $A = \mathcal{O}_K$ and $B = \mathcal{O}_L$. Assume that $\alpha \leq \beta$. Then $a_\beta \mid a_\alpha$, so that $\frac{a_\alpha}{a_\beta}, \left(\frac{a_\alpha}{a_\beta} \right)^{n-1} \in \mathcal{O}_K$. From (1) and (43) it thus follows that $b_\alpha \in (b_\beta)$ and $h'_\alpha(b_\alpha) \in (h'_\beta(b_\beta))$. This shows that \mathcal{O}_L -ideals $(b_\alpha)_{\alpha \in S}$ and $(h'_\alpha(a_\alpha))_{\alpha \in S}$ form increasing chains.

Using (43) with $\beta = \gamma$, we obtain:

$$(44) \quad V = (h'_\alpha(b_\alpha) \mid \alpha \in S) = \left(\left(\frac{a_\alpha}{a_\gamma} \right)^{n-1} h'_\gamma(b_\gamma) \mid \alpha \in S \right) = \frac{h'_\gamma(b_\gamma)}{a_\gamma^{n-1}} U^{n-1}.$$

This proves (3).

4.2. The ideal V of differents.

Under the assumptions of Theorem 1.1, with $(L|K, v)$ a valued field extension satisfying condition (8) and $A = \mathcal{O}_K$ and $B = \mathcal{O}_L$, we will now show that V is the \mathcal{O}_L -ideal generated by the differents of the generators b_α of \mathcal{O}_L over \mathcal{O}_K . Take any generator $b \in \mathcal{O}_L$ of $L|K$ and let h be its minimal polynomial over K . Denote the different of b by $\delta(b) := h'(b)$. For $i \geq 1$ we have

$$b^i - \sigma b^i = b^i - (\sigma b)^i = b^i - (b + (\sigma b - b))^i = \sum_{j=0}^{i-1} \binom{i}{j} b^j (\sigma b - b)^{i-j}.$$

Since the extension is unbranched, we have $v\sigma b = vb$, whence $v(b - \sigma b) \geq vb \geq 0$. Consequently,

$$v(b^i - \sigma b^i) \geq v(\sigma b - b) = v(b - \sigma b).$$

Every $b \in \mathcal{O}_K[b_\alpha] \setminus \mathcal{O}_K$ is of the form

$$b = \sum_{i=0}^{n-1} c_i b_\alpha^i$$

with $c_i \in \mathcal{O}_K$. Therefore,

$$\begin{aligned} v\delta(b) &= v \prod_{\text{id} \neq \sigma \in G} (b - \sigma b) = \sum_{\text{id} \neq \sigma \in G} v \left(\sum_{i=0}^{n-1} c_i b_\alpha^i - \sigma \sum_{i=0}^{n-1} c_i (b_\alpha)^i \right) \\ &= \sum_{\text{id} \neq \sigma \in G} v \sum_{i=1}^{n-1} c_i (b_\alpha^i - \sigma b_\alpha^i). \end{aligned}$$

For $1 \leq i \leq n-1$, we have

$$v c_i (b_\alpha^i - \sigma b_\alpha^i) \geq v (b_\alpha^i - \sigma b_\alpha^i) \geq v (b_\alpha - \sigma b_\alpha),$$

showing that

$$v \sum_{i=1}^{n-1} c_i (b_\alpha^i - \sigma b_\alpha^i) \geq v (b_\alpha - \sigma b_\alpha).$$

Hence,

$$v\delta(b) \geq \sum_{\text{id} \neq \sigma \in G} v (b_\alpha - \sigma b_\alpha) = v h'_\alpha(b_\alpha).$$

We use this to conclude:

Proposition 4.1. *Under the above assumptions, we have:*

$$(45) \quad V = (\delta(b) \mid b \in \mathcal{O}_L \setminus \mathcal{O}_K).$$

Proof. Since \mathcal{O}_L is the union of the chain of rings $\mathcal{O}_K[b_\alpha]$, $\alpha \in S$, we have

$$(\delta(b) \mid b \in \mathcal{O}_L \setminus \mathcal{O}_K) = \bigcup_{\alpha \in S} (\delta(b) \mid b \in \mathcal{O}_K[b_\alpha] \setminus \mathcal{O}_K) = \bigcup_{\alpha \in S} (h'_\alpha(b_\alpha)) = V.$$

□

4.3. The case of Artin-Schreier defect extensions.

Theorem 4.2. *Let $(L|K, v)$ be an Artin-Schreier defect extension with ramification ideal $I_\mathcal{E}$. Then there is an \mathcal{O}_L -module isomorphism*

$$\Omega_{\mathcal{O}_L/\mathcal{O}_K} \cong I_\mathcal{E}/I_\mathcal{E}^p.$$

Proof. Let ϑ be an Artin-Schreier generator of $L|K$. By Theorem 3.4, the \mathcal{O}_K -algebras $\mathcal{O}_K[\vartheta_c]$, with $c \in K$, form a chain of subrings of L whose union is \mathcal{O}_L . For $c \in K$, $\vartheta_c = t_c(\vartheta - c)$, where $t_c \in \mathcal{O}_K$ is such that $v\vartheta_c = 0$. For $v(\vartheta - c_1) < v(\vartheta - c_2)$, by (28) we have that

$$\vartheta_{c_1} = \frac{t_{c_1}}{t_{c_2}} \vartheta_{c_2} + t_{c_1}(c_2 - c_1)$$

with $\frac{t_{c_1}}{t_{c_2}}, t_{c_1}(c_2 - c_1) \in \mathcal{O}_K$. We will apply Theorem 1.1 with $A = \mathcal{O}_K$, $B = \mathcal{O}_L$, and $S = v(\vartheta - K)$. For each $\alpha \in S$ we choose $c_\alpha \in K$ such that $v(\vartheta - c_\alpha) = \alpha$ and set $b_\alpha = \vartheta_{c_\alpha}$. We set $a_\alpha = t_{c_\alpha}$ and $c_{\alpha,\beta} = t_{c_\alpha}(c_\beta - c_\alpha)$. We denote by h_α the minimal polynomial of $b_\alpha = \vartheta_{c_\alpha}$ over K . Thus in the notation of Lemma 3.6, $h_\alpha = g_{c_\alpha}$ so that $h'_\alpha(b_\alpha) = g'_{c_\alpha}(\vartheta_{c_\alpha}) = -t_{c_\alpha}^{p-1}$ by equation (38). Hence for every $\alpha \in S$, $h'_\alpha(b_\alpha)/a_\alpha^{p-1} = -1$. Our theorem now follows from equation (3) of Theorem 1.1, together with Lemma 3.6 which shows that the \mathcal{O}_L -ideal $U = (t_\alpha \mid \alpha \in S)$ is equal to $I_\mathcal{E}$. \square

4.4. The case of Kummer defect extensions of degree p .

Theorem 4.3. *Assume that $\text{char } K = 0$ and let $(L|K, v)$ be a Kummer defect extension of degree p with ramification ideal $I_\mathcal{E}$. Then there is an \mathcal{O}_L -module isomorphism*

$$\Omega_{\mathcal{O}_L|\mathcal{O}_K} \cong I_\mathcal{E}/I_\mathcal{E}^p.$$

Proof. Let η be a Kummer generator of $L|K$ which is a 1-unit. We use the notation from Section 3.2. By Corollary 3.5, the \mathcal{O}_K -algebras $\mathcal{O}_K[\eta_{<c>}]$ with c a 1-unit in K form a chain of subrings of L whose union is \mathcal{O}_L . For $v(\eta - c_1) < v(\eta - c_2)$, by (33) we have that

$$\eta_{<c_1>} = \frac{T_{c_1}}{T_{c_2}}\eta_{<c_2>} + T_{c_1}\chi(c_2 - c_1)$$

with $\frac{T_{c_1}}{T_{c_2}}, T_{c_1}\chi(c_2 - c_1) \in \mathcal{O}_K$. We will apply Theorem 1.1 with $A = \mathcal{O}_K$, $B = \mathcal{O}_L$, and $S = v(\eta - K)$. For each $\alpha \in S$ we choose $c_\alpha \in K$ such that $v(\eta - c_\alpha) = \alpha$. Let $b_\alpha = \eta_{<c_\alpha>}$, $a_\alpha = T_{c_\alpha}$, and $c_{\alpha,\beta} = T_{c_\alpha}\chi(c_\beta - c_\alpha)$. We denote by h_α the minimal polynomial of $b_\alpha = \eta_{<c_\alpha>}$ over K . Thus in the notation of Lemma 3.7, $h_\alpha = f_{c_\alpha}$ so that by equation (40), $h'_\alpha(b_\alpha) = f'_{c_\alpha}(\eta_{<c_\alpha>}) = T_{c_\alpha}^{p-1}u$ with u a unit in \mathcal{O}_L . Hence for every $\alpha \in S$, $h'_\alpha(b_\alpha)/a_\alpha^{p-1} = u$. Our theorem now follows from equation (3) of Theorem 1.1, together with Lemma 3.7, which shows that the \mathcal{O}_L -ideal $U = (T_\alpha \mid \alpha \in S)$ is equal to $I_\mathcal{E}$. \square

4.5. The valuation ring $\mathcal{O}(I)$ and the annihilator.

Take an \mathcal{O}_L -ideal I . We define $\mathcal{O}(I)$ to be the largest valuation ring \mathcal{O} containing \mathcal{O}_L such that I is an \mathcal{O} -ideal. Further, we define $H(I)$ to be the largest convex subgroup H of vL such that $vI + H = vI$. In the terminology of [18, 19], this is called the **invariance group** of vI , denoted by $\mathcal{G}(vI)$. We infer from [19, Theorem 3.6]:

Proposition 4.4. *For every \mathcal{O}_L -ideal I ,*

$$(46) \quad \mathcal{O}(I) = \mathcal{O}_{v_{H(I)}}.$$

Accordingly, we set $\mathcal{M}(I) := \mathcal{M}_{v_{H(I)}}$. We will use the following facts:

Proposition 4.5. *Take an \mathcal{O}_L -ideal U and set $V = bU^{n-1}$, where $b \in \mathcal{O}_L$.*

- 1) *We have $H(U) = H(V)$, $\mathcal{O}(U) = \mathcal{O}(V)$ and $\mathcal{M}(U) = \mathcal{M}(V)$.*
- 2) *If there is $a \in K$ such that $v_{H(V)}V$ has infimum $v_{H(V)}a$ in $v_{H(V)}K$ but $v_{H(V)}V$ does not contain this infimum, then*

$$(47) \quad \text{ann } U/UV = a\mathcal{O}(V),$$

which properly contains V . In all other cases, $\text{ann } U/UV = V$.

In all cases, $\mathcal{M}(V) \text{ann } U/UV \subseteq V$.

3) If \mathcal{M}_L annihilates U/UV , then $U/UV = 0$, except if \mathcal{M}_L is principal and either $U = V = \mathcal{M}_L$ (which holds if and only if $n = 2$ and $b \in \mathcal{O}_L^\times$), or $U = \mathcal{O}_L$ and $V = \mathcal{M}_L = (b)$ (which holds if and only if $n = 1$ and $b \notin \mathcal{O}_L^\times$). In these exceptional cases, $\mathcal{M}_L = \text{ann } U/UV$.

Proof. 1) follows from [19, Lemmas 2.14 and 3.7]. 2) and 3) follow from [19, Proposition 3.19]. \square

In Theorems 4.2 and 4.3 we proved that in the case of $\mathcal{E} = (L|K, v)$ an Artin-Schreier or a Kummer defect extension of degree p with ramification ideal $I_{\mathcal{E}}$ we have $\Omega_{\mathcal{O}_L/\mathcal{O}_K} \cong I_{\mathcal{E}}/I_{\mathcal{E}}^p$. Therefore, we will now assume that the extension $\mathcal{E} = (L|K, v)$ satisfies (8) and compute the corresponding annihilator. We observe:

Lemma 4.6. *Assume that the extension $\mathcal{E} = (L|K, v)$ satisfies (8) and has independent defect. Then*

$$H(vI_{\mathcal{E}}) = H_{\mathcal{E}}.$$

Proof. We have $vI_{\mathcal{E}} = \Sigma_{\mathcal{E}} = \{\alpha \in vL \mid \alpha > H_{\mathcal{E}}\}$. In [19], this set is denoted by $H_{\mathcal{E}}^+$, and in [19, part 5) of Lemma 2.14] (where we take $\alpha = 0$) it is shown that for each convex subgroup H the invariance group of H^+ equals H . \square

Analogously, we define $H_{\mathcal{E}} := H(vI_{\mathcal{E}})$ in case \mathcal{E} has dependent defect. Then we obtain from Proposition 4.4 that

$$(48) \quad \mathcal{O}(I_{\mathcal{E}}) = \mathcal{O}_{vH_{\mathcal{E}}}$$

holds for all extensions $\mathcal{E} = (L|K, v)$ satisfying (8). Applying Proposition 4.5, we find:

Proposition 4.7. *Assume that the extension $\mathcal{E} = (L|K, v)$ satisfies (8).*

1) *We have $H(I_{\mathcal{E}}) = H(I_{\mathcal{E}}^{p-1}) = H(I_{\mathcal{E}}^p)$, $\mathcal{O}(I_{\mathcal{E}}) = \mathcal{O}(I_{\mathcal{E}}^{p-1}) = \mathcal{O}(I_{\mathcal{E}}^p)$ and $\mathcal{M}(I_{\mathcal{E}}) = \mathcal{M}(I_{\mathcal{E}}^{p-1}) = \mathcal{M}(I_{\mathcal{E}}^p)$.*

2) *If there is $a \in K$ such that $v_{H_{\mathcal{E}}}I_{\mathcal{E}}^{p-1}$ has infimum $v_{H_{\mathcal{E}}}a$ in $v_{H_{\mathcal{E}}}K$ but $v_{H_{\mathcal{E}}}I_{\mathcal{E}}^p$ does not contain this infimum, then*

$$(49) \quad \text{ann } I_{\mathcal{E}}/I_{\mathcal{E}}^p = a\mathcal{O}(I_{\mathcal{E}}),$$

which properly contains $I_{\mathcal{E}}^{p-1}$. In all other cases, $\text{ann } I_{\mathcal{E}}/I_{\mathcal{E}}^p = I_{\mathcal{E}}^{p-1}$.

In all cases, $\mathcal{M}(I_{\mathcal{E}}) \text{ann } I_{\mathcal{E}}/I_{\mathcal{E}}^p \subseteq I_{\mathcal{E}}^{p-1}$.

3) *If \mathcal{M}_L annihilates $\Omega_{\mathcal{O}_L/\mathcal{O}_K}$, then $\Omega_{\mathcal{O}_L/\mathcal{O}_K} = 0$ and $\text{ann } \Omega_{\mathcal{O}_L/\mathcal{O}_K} = \mathcal{O}_L$.*

Proof. Part 1) follows from [19, Lemmas 2.14 and 3.7]. Part 2) follows from part 2) of Proposition 4.5 where $U = I_{\mathcal{E}}$ and $V = I_{\mathcal{E}}^{p-1}$, and using that $\mathcal{O}(I_{\mathcal{E}}) = \mathcal{O}(I_{\mathcal{E}}^{p-1})$. Part 3) follows from Theorems 4.2 and 4.3 and part 3) of Proposition 4.5 since under our assumptions on \mathcal{E} , $I_{\mathcal{E}}^{p-1}$ cannot be principal. \square

5. THE TRACE OF VALUATION RINGS AND DIFFERENTS

5.1. Defect extensions of prime degree.

In this section we will consider the trace on an extension $\mathcal{E} = (L|K, v)$ that satisfies (8). If $L|K$ is an Artin-Schreier extension, then we write $L = K(\vartheta)$ where ϑ is an Artin-Schreier generator. If $\text{char } K = 0$ and $L|K$ is a Kummer extension, then we write $L = K(\eta)$ where η is a Kummer generator, that is, $\eta^p \in K$; as explained at the beginning of Section 2.8, we can assume that η is a 1-unit.

Take $a \in L \setminus K$. The proof of the following fact can be found in [14, Section 6.3].

Lemma 5.1. *Take a separable field extension $K(a)|K$ of degree n and let $f(X) \in K[X]$ be the minimal polynomial of a over K . Then*

$$(50) \quad \text{Tr}_{K(a)|K} \left(\frac{a^m}{f'(a)} \right) = \begin{cases} 0 & \text{if } 1 \leq m \leq n-2 \\ 1 & \text{if } m = n-1. \end{cases}$$

□

For arbitrary $b, c \in K$, we note:

$$(51) \quad b(a-c)^{p-1} \in \mathcal{M}_{K(a)} \iff vb > -(p-1)v(a-c).$$

First we consider Artin-Schreier extensions. By Lemma 5.1 and equation (35),

$$(52) \quad \text{Tr}_{K(\vartheta)|K} ((\vartheta - c)^i) = \begin{cases} 0 & \text{if } 1 \leq i \leq p-2 \\ -1 & \text{if } i = p-1 \end{cases}$$

for arbitrary $c \in K$ since $\vartheta - c$ is also an Artin-Schreier generator. In particular,

$$\text{Tr}_{K(\vartheta)|K} (b(\vartheta - c)^{p-1}) = -b$$

for all $b \in K$. By (51) it follows that

$$(53) \quad \begin{aligned} \text{Tr}_{K(\vartheta)|K} (\mathcal{M}_{K(\vartheta)}) &\supseteq \{b \in K \mid vb > -(p-1)v(\vartheta - c) \text{ for some } c \in K\} \\ &= \{b \in K \mid vb \geq -(p-1)v(\vartheta - c) \text{ for some } c \in K\} \\ &= \{b \in K \mid vb \in -(p-1)v(\vartheta - K)\} \\ &= \{b \in K \mid vb \in (p-1)\Sigma_{\mathcal{E}}\}, \end{aligned}$$

where the first equality follows from the fact that $-(p-1)v(\vartheta - c)$ has no smallest element, and the last equality follows from equation (23) of Theorem 2.8.

Now we consider Kummer extensions. Since $(\eta^i)^p \in K$, we have that

$$\text{Tr}_{K(\eta)|K} (\eta^i) = 0$$

for $1 \leq i \leq p-1$. For $c \in K$ and $0 \leq j \leq p-1$, we compute:

$$(\eta - c)^j = \sum_{i=1}^j \binom{j}{i} \eta^i (-c)^{j-i} + (-c)^j.$$

Thus for every $b \in K$,

$$(54) \quad \text{Tr}_{K(\eta)|K} (b(\eta - c)^j) = pb(-c)^j.$$

By (51), $b(\eta - c)^{p-1} \in \mathcal{M}_{K(\eta)}$ holds if and only if $vb > -(p-1)v(\eta - c)$; the latter remains true if we make $v(\eta - c)$ even larger. Since η is a 1-unit, there is $c \in K$ such that $v(\eta - c) > 0$, which implies that $vc = 0$. Hence we may choose $c \in K$

with $vb > -(p-1)v(\eta - c)$ and $vc = 0$. Applying (54) with $j = p-1$, we find that $\text{Tr}_{K(\eta)|K}(b(-c)^{-(p-1)}(\eta - c)^{p-1}) = pb$. We obtain:

$$\begin{aligned}
 (55) \quad \text{Tr}_{K(\eta)|K}(\mathcal{M}_{K(\eta)}) &\supseteq \{pb \mid b \in K, vb > -(p-1)v(\eta - c) \text{ for some } c \in K\} \\
 &= \{b \in K \mid vb \geq vp - (p-1)v(\eta - c) \text{ for some } c \in K\} \\
 &= \left\{ b \in K \mid vb \in (p-1) \left(\frac{1}{p-1}vp - v(\eta - K) \right) \right\} \\
 &= \{b \in K \mid vb \in (p-1)\Sigma_{\mathcal{E}}\},
 \end{aligned}$$

where the first equality follows from the fact that $-(p-1)v(\vartheta - c)$ has no smallest element, and the last equality follows from equation (24) of Theorem 2.8.

In order to prove the opposite inclusions in (53) and (55), we have to find out enough information about the elements $g(a) \in K(a)$ that lie in $\mathcal{M}_{K(a)}$. Using the Taylor expansion, we write

$$g(a) = \sum_{i=0}^{p-1} \partial_i g(c)(a-c)^i.$$

By Lemma 2.4 there is $c \in K$ such that among the values $v\partial_i g(c)(a-c)^i$, $0 \leq i \leq p-1$, there is precisely one of minimal value, and the same holds for all $c' \in K$ with $v(a-c') \geq v(a-c)$. In particular, we may assume that $v(a-c) > va$. For all such c , we have:

$$vg(a) = \min_{0 \leq i \leq p-1} v\partial_i g(c)(a-c)^i.$$

Hence for $g(a)$ to lie in $\mathcal{M}_{K(a)}$ it is necessary that $v\partial_i g(c)(a-c)^i > 0$, or equivalently,

$$(56) \quad v\partial_i g(c) > -iv(a-c)$$

for $0 \leq i \leq p-1$ and $c \in K$ with $v(a-c) > va$.

In the Artin-Schreier extension case, for $g(\vartheta) \in \mathcal{M}_{K(\vartheta)}$ and $c \in K$ with $v(a-c) > va$, using (52) we find:

$$\text{Tr}_{K(\vartheta)|K}(g(\vartheta)) = \sum_{i=0}^{p-1} \text{Tr}_{K(\vartheta)|K}(\partial_i g(c)(\vartheta - c)^i) = -\partial_{p-1}g(c).$$

Since $\partial_{p-1}g(c) > -(p-1)v(\vartheta - c)$ by (56), this proves the desired equality in (53).

In the Kummer extension case, for $g(\eta) \in \mathcal{M}_{K(\eta)}$ and $c \in K$ with $v(a-c) > va$, using (54) we find:

$$\text{Tr}_{K(\eta)|K}(g(\eta)) = \sum_{j=0}^{p-1} \text{Tr}_{K(\eta)|K}(\partial_j g(c)(\eta - c)^j) = p \sum_{j=0}^{p-1} \partial_j g(c)(-c)^j.$$

As we assume that $v(\eta - c) > 0$, we have that $vc = 0$ and

$$-iv(\eta - c) \geq -(p-1)v(\eta - c) \quad \text{for } 0 \leq i \leq p-1.$$

Hence by (56), $v \sum_{i=0}^{p-1} \partial_i g(c)(-c)^i \geq -(p-1)v(\eta - c)$. This proves the desired equality in (55).

Remark 5.2. In the case of a Kummer extension, it follows from (55) that $p\mathcal{O}_K \subseteq \text{Tr}_{K(\eta)|K}(\mathcal{M}_{K(\eta)})$ since $vp = vp - (p-1)v\eta$ as η is a unit.

5.2. Proof of Theorem 1.5.

We have just proved the second equality of (9) in the previous section; the third equality follows from part 1) of Lemma 2.7. The first equality is seen as follows. Take any $a \in \mathcal{O}_L$. As the defect extension $(L|K, v)$ is immediate, there is some $b \in \mathcal{O}_K$ such that $a - b \in \mathcal{M}_L$. Then $\text{Tr}_{L|K}(a) = \text{Tr}_{L|K}(b) + \text{Tr}_{L|K}(a - b)$ with $\text{Tr}_{L|K}(a - b) \in \text{Tr}_{L|K}(\mathcal{M}_L)$. If $\text{char } L = p$, then $\text{Tr}_{L|K}(b) = 0$. If $\text{char } L = 0$, then $\text{Tr}_{L|K}(b) = pb \in p\mathcal{O}_K$, which is contained in $\text{Tr}_{L|K}(\mathcal{M}_L)$ as stated in Remark 5.2.

Assume that the extension \mathcal{E} has independent defect. Then by part 1) of Proposition 2.10, $I_{\mathcal{E}}^p = I_{\mathcal{E}}$ as well as $(I_{\mathcal{E}} \cap K)^p = I_{\mathcal{E}} \cap K$. This implies that $I_{\mathcal{E}}^{p-1} = I_{\mathcal{E}}$ and $(I_{\mathcal{E}} \cap K)^{p-1} = I_{\mathcal{E}} \cap K$. On the other hand, we know from the equivalence of statements a) and b) in Theorem 1.4 that $I_{\mathcal{E}} = \mathcal{M}_{v_H}$, where H is a strongly convex subgroup of vL , and that H is equal to the convex subgroup $H_{\mathcal{E}}$ appearing in the definition of independent defect. Now the second equation of (10) follows from (9) as $(I_{\mathcal{E}} \cap K)^{p-1} = I_{\mathcal{E}} \cap K = \mathcal{M}_{v_H} \cap K$. The third equation of (10) holds since $\mathcal{M}_{v_H} = (a \in L \mid va > H)$.

Now assume that (10) holds for some strongly convex subgroup H of $vL = vK$. Set $\Sigma := vL \setminus H = vK \setminus H$. From (9) we now obtain:

$$(b \in K \mid vb \in (p-1)\Sigma_{\mathcal{E}}) = (b \in K \mid vb \in \Sigma).$$

On the one hand, we have

$$(b \in K \mid vb \in (p-1)\Sigma_{\mathcal{E}}) = (b \in K \mid vb \in ((p-1)\Sigma_{\mathcal{E}})^-),$$

and on the other,

$$(b \in K \mid vb \in \Sigma) = (b \in K \mid vb \in ((p-1)\Sigma)^-)$$

by Lemma 2.1. This implies that $((p-1)\Sigma_{\mathcal{E}})^- = ((p-1)\Sigma)^-$. Suppose that $\Sigma_{\mathcal{E}} \neq \Sigma$. If there is $\alpha \in \Sigma_{\mathcal{E}} \setminus \Sigma$, then $\alpha < \Sigma$, whence $(p-1)\alpha < (p-1)\Sigma$ and therefore, $((p-1)\Sigma_{\mathcal{E}})^- \not\supseteq ((p-1)\Sigma)^-$, contradiction. Symmetrically, we obtain a contradiction if there is $\alpha \in \Sigma \setminus \Sigma_{\mathcal{E}}$. This shows that $\Sigma = \Sigma_{\mathcal{E}}$, and we thus obtain that \mathcal{E} has independent defect and that $H = H_{\mathcal{E}}$.

The last statement of the theorem holds since $vp \notin H_{\mathcal{E}} = H$ by part 2) of Proposition 2.10. If $(vK)_{vp}$ is archimedean, this forces $H = \{0\}$. \square

5.3. Differents.

In this section, we compute the differents $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K)$ in the case where $vL = vK$. We will employ the following auxiliary results.

Lemma 5.3. *Take any \mathcal{O}_L -fractional ideal I .*

1) *If vI has an infimum α in vL , then $\mathcal{O}_L :_L I = (b \in L \mid vb \geq -\alpha)$. If vI has no infimum in vL , then $\mathcal{O}_L :_L I = (b \in L \mid vb > -vI)$, which is not a principal ideal, and $v(\mathcal{O}_L :_L I)$ has no infimum.*

2) *If vI has an infimum α in vL , then $\mathcal{O}_L :_L (\mathcal{O}_L :_L I) = (b \in L \mid vb \geq \alpha)$. If vI has no infimum in vL , then $\mathcal{O}_L :_L (\mathcal{O}_L :_L I) = I$.*

Proof. 1): We compute:

$$\begin{aligned} \mathcal{O}_L :_L I &= (b \in L \mid bI \subseteq \mathcal{O}_L) = (b \in L \mid vb + \beta \geq 0 \text{ for all } \beta \in vI) \\ &= (b \in L \mid vb \geq -vI). \end{aligned}$$

Since α is an infimum of vI if and only if $-\alpha$ is a supremum of $-vI$, this yields our assertions.

2): This follows by applying part 1) twice. \square

Denote by $\mathcal{T}(\mathcal{O}_L|\mathcal{O}_K)$ the \mathcal{O}_L -ideal generated by $\text{Tr}_{L|K}(\mathcal{O}_L)$. We use the previous lemma to show:

Theorem 5.4. *Assume that $(L|K, v)$ is a finite unbranched extension with $vL = vK$. Then we have:*

1) $\mathcal{C}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{O}_L :_L \mathcal{T}(\mathcal{O}_L|\mathcal{O}_K)$ and $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{O}_L :_L (\mathcal{O}_L :_L \mathcal{T}(\mathcal{O}_L|\mathcal{O}_K))$.

2) If $v\text{Tr}_{L|K}(\mathcal{O}_L)$ has an infimum α in vK , then $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = (b \in L \mid vb \geq \alpha)$ and if $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) \neq \mathcal{T}(\mathcal{O}_L|\mathcal{O}_K)$, then

$$(57) \quad \mathcal{M}_L \mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{T}(\mathcal{O}_L|\mathcal{O}_K).$$

If $v\text{Tr}_{L|K}(\mathcal{O}_L)$ has no infimum in vK , then $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{T}(\mathcal{O}_L|\mathcal{O}_K)$, which is not principal.

Proof. 1): Take any $z \in L$. Since $vL = vK$, there is $z_1 \in K$ such that $vz_1 = vz$, so that $z = z_1 z_2$ with z_2 a unit in \mathcal{O}_L . Then

$$\text{Tr}_{L|K}(z\mathcal{O}_L) = z_1 \text{Tr}_{L|K}(z_2\mathcal{O}_L) = z_1 \text{Tr}_{L|K}(\mathcal{O}_L).$$

Observe that $z_1 \text{Tr}_{L|K}(\mathcal{O}_L) \subseteq \mathcal{O}_K$ if and only if $z\mathcal{T}(\mathcal{O}_L|\mathcal{O}_K) \subseteq \mathcal{O}_L$. This proves the assertions of part 1).

2): A part of the assertions follow from part 1) together with part 2) of Lemma 5.3; it only remains to prove (57) in case $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K)$ properly contains $\mathcal{T}(\mathcal{O}_L|\mathcal{O}_K)$. In this case, $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = (b \in L \mid vb \geq \alpha)$ and $\mathcal{T}(\mathcal{O}_L|\mathcal{O}_K) = (b \in L \mid vb > \alpha) = v\mathcal{M}_L + (b \in L \mid vb \geq \alpha)$, which proves (57). \square

We will also need the following simple observation.

Lemma 5.5. *Take an extension $(L|K, v)$ of valued fields with $vL = vK$, an \mathcal{O}_L -ideal I , and $n \in \mathbb{N}_{>0}$. Then the \mathcal{O}_L -ideal J generated by $(I \cap K)^n$ equals I^n .*

Proof. The inclusion $J \subseteq I^n$ is obvious. Now take any $b \in I$. Since $vL = vK$, there is $c \in K$ such that $vc = vb$, which implies that $c \in I \cap K$. Now $b^n = (b/c)^n c^n$ with $(b/c)^n \in \mathcal{O}_L$ and $c^n \in I \cap K$. Therefore, $b^n \in J$, which proves the reverse inclusion. \square

Proof of Theorem 1.6:

1): Lemmas 3.6 and 3.7 together with Proposition 4.1 show that i) and ii) are equal. Proposition 4.1 shows that ii) and iii) are equal. From Theorem 1.5 we know that $\text{Tr}_{L|K}(\mathcal{O}_L) = (I_{\mathcal{E}} \cap K)^{p-1}$, whence $\mathcal{T}(\mathcal{O}_L|\mathcal{O}_K) = I_{\mathcal{E}}^{p-1}$ by Lemma 5.5. This shows that i) and iv) are equal.

2): If $v\mathcal{T}(\mathcal{O}_L|\mathcal{O}_K) = vI_{\mathcal{E}}^{p-1}$ has no infimum in $vK = vL$, then from Theorem 5.4 we obtain that $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{T}(\mathcal{O}_L|\mathcal{O}_K) = I_{\mathcal{E}}^{p-1}$. As the extension $(L|K, v)$ is immediate, $vI_{\mathcal{E}}^{p-1}$ has no smallest element. Hence if $vI_{\mathcal{E}}^{p-1}$ has an infimum α , it does not contain it, and again by Theorem 5.4, $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = (b \in L \mid vb \geq \alpha) \supsetneq I_{\mathcal{E}}^{p-1}$ and $\mathcal{M}_L \mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{T}(\mathcal{O}_L|\mathcal{O}_K) = I_{\mathcal{E}}^{p-1}$. Note that if $\alpha = va$ for $a \in L$, then $(b \in L \mid vb \geq \alpha) = a\mathcal{O}_L$.

3): Assume that (K, v) has rank 1. Then the only proper convex subgroup of vL is $\{0\}$. Thus Proposition 4.7 together with Theorems 4.2 and 4.3 shows that $\text{ann } \Omega_{\mathcal{O}_L|\mathcal{O}_K} = I_{\mathcal{E}}^{p-1}$ if $vI_{\mathcal{E}}^{p-1}$ has no infimum in vL , and $\text{ann } \Omega_{\mathcal{O}_L|\mathcal{O}_K} = \{b \in L \mid vb \geq \alpha\}$ if $vI_{\mathcal{E}}^{p-1}$ has infimum α . From what we have already shown, it follows that $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = \text{ann } \Omega_{\mathcal{O}_L|\mathcal{O}_K}$.

4): Assume that the extension \mathcal{E} has independent defect; then $\Omega_{\mathcal{O}_L|\mathcal{O}_K} = 0$ by Theorem 1.4 and therefore, $\text{ann } \Omega_{\mathcal{O}_L|\mathcal{O}_K} = \mathcal{O}_L$. If $H_{\mathcal{E}} = \{0\}$, then $\text{Tr}_{L|K}(\mathcal{O}_L) = \text{Tr}_{L|K}(\mathcal{M}_L) = \mathcal{M}_K$ by Theorem 1.5. Since $H_{\mathcal{E}} = \{0\}$ is a strongly convex subgroup of $vL = vK$, we know that vK has no minimal positive element. So $v\mathcal{M}_K$ has infimum 0 and it follows from part 2) of Theorem 5.4 that $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{O}_L$.

Now assume that $H_{\mathcal{E}} \neq \{0\}$. Since $\mathcal{M}_{v_{H_{\mathcal{E}}}} \cap K = \mathcal{M}_{v_{H_{\mathcal{E}}}|K}$, Lemma 5.5 shows that the \mathcal{O}_L -ideal generated by $\mathcal{M}_{v_{H_{\mathcal{E}}}} \cap K$ is $\mathcal{M}_{v_{H_{\mathcal{E}}}}$. By Theorem 1.5, $\text{Tr}_{L|K}(\mathcal{O}_L) = \text{Tr}_{L|K}(\mathcal{M}_L) = \mathcal{M}_{v_{H_{\mathcal{E}}}} \cap K$, so $\mathcal{T}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{M}_{v_{H_{\mathcal{E}}}}$. As $v(\mathcal{M}_{v_{H_{\mathcal{E}}}} \cap K) = v\mathcal{M}_{v_{H_{\mathcal{E}}}}$ has no infimum in vK , part 2) of Theorem 5.4 shows that $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{T}(\mathcal{O}_L|\mathcal{O}_K) = \mathcal{M}_{v_{H_{\mathcal{E}}}} \subsetneq \mathcal{O}_L = \text{ann } \Omega_{\mathcal{O}_L|\mathcal{O}_K}$ in this case.

Finally, assume that $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K)$ is equal to \mathcal{O}_L or to \mathcal{M}_{v_H} for a strongly convex nontrivial subgroup H of vL . If the former holds, then $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K)$ is not equal to $I_{\mathcal{E}}^{p-1}$ since $vI_{\mathcal{E}} = \Sigma_{\mathcal{E}}$ and hence also $I_{\mathcal{E}}^{p-1}$ have no smallest element. Hence from the proof of part 2) it follows that 0 is the infimum of $vI_{\mathcal{E}}^{p-1}$. This implies that $I_{\mathcal{E}} = \mathcal{M}_L = \mathcal{M}_{v_H}$ where $H = \{0\}$. Since $v\mathcal{M}_L = vI_{\mathcal{E}}$ has no smallest element, it follows that $\{0\}$ is a strongly convex subgroup of vL .

If the latter holds, then $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K)$ is not principal and therefore by part 2), $\mathcal{D}(\mathcal{O}_L|\mathcal{O}_K)$ can only be equal to $I_{\mathcal{E}}^{p-1}$. This implies that $I_{\mathcal{E}} = \mathcal{M}_{v_H}$ for some strongly convex subgroup of vL . In both cases, it follows from the equivalence a) \Leftrightarrow b) of Theorem 1.4 that \mathcal{E} has independent defect. \square

6. PROOF OF THEOREMS 1.7 AND 1.8

Take a defect extension $\mathcal{E} = (L|K, v)$ of degree p . Then by equation (13), the extension is unbranched and immediate, so in particular, $vL = vK$.

We first prove Theorem 1.7, so we assume that \mathcal{E} is an Artin-Schreier extension generated by $\vartheta \in L$ with $\vartheta^p - \vartheta = a \in K$. By Corollary 2.9, $v(\vartheta - c) < 0$ for all $c \in K$. It follows that $v(\vartheta^p - c^p) = v(\vartheta - c)^p = pv(\vartheta - c) < v(\vartheta - c)$ and thus,

$$(58) \quad v(a - \wp(c)) = v(\vartheta^p - \vartheta - c^p + c) = \min\{v(\vartheta^p - c^p), v(\vartheta - c)\} = pv(\vartheta - c).$$

This proves equation (11).

The equivalence a) \Leftrightarrow b) follows from the definition of independent defect together with equation (23) of Theorem 2.8 which says that $\Sigma_{\mathcal{E}} = -v(\vartheta - K)$. The equivalences b) \Leftrightarrow c) and e) \Leftrightarrow f) follow from equation (11). The equivalence c) \Leftrightarrow d) is trivial.

Now assume that vK is p -divisible. Then the equivalence a) \Leftrightarrow e) follows from part 1) of Proposition 2.10 together with the equation $\Sigma_{\mathcal{E}} = -v(\vartheta - K)$.

Finally, assume that (K, v) has rank 1. Then the only proper convex subgroup of vK is $H = \{0\}$, so that condition d) is equivalent to condition g).

Now we prove Theorem 1.8, so we assume that \mathcal{E} is a Kummer extension. As explained at the beginning of Section 2.8, it is generated by a 1-unit $\eta \in L$ with $\eta^p = a \in K$. By Corollary 2.9, $v(\eta - c) < \frac{1}{p-1}vp$ for all $c \in K$.

The following is part of Lemma 2.18 of [20]:

Lemma 6.1. *Take $\eta \in \tilde{K}$ such that $\eta^p \in K$ and $v\eta = 0$. Then for $c \in K$ such that $v(\eta - c) > 0$, $v(\eta - c) < \frac{1}{p-1}vp$ holds if and only if $v(\eta^p - c^p) < \frac{p}{p-1}vp$, and if this is the case, then $v(\eta^p - c^p) = pv(\eta - c)$.*

In order to prove equation (12), we have to discuss the remaining case of $v(\eta - c) \leq 0$. If $v(\eta - c) = 0$, then $vc \geq 0$ and c is not a 1-unit. Hence c^p is not a 1-unit while $a = \eta^p$ is, whence $v(a - c^p) = 0 = pv(\eta - c)$. If $v(\eta - c) < 0$, then $vc < 0$ and $v(\eta - c) = vc$. It follows that $vc^p < 0 = v\eta^p$, so that $v(\eta^p - c^p) = vc^p = pvc = pv(\eta - c)$. This completes the proof of equation (12).

The equivalence a) \Leftrightarrow b) follows from the definition of independent defect together with equation (24) of Theorem 2.8, which says that $\Sigma_{\mathcal{E}} = \frac{1}{p-1}vp - v(\vartheta - K)$. The equivalences b) \Leftrightarrow c) and e) \Leftrightarrow f) follow from equation (12) and the fact that $\alpha < H$ if and only if $p\alpha < H$ since H is a convex subgroup of vK . The equivalence c) \Leftrightarrow d) is trivial.

Now assume that vK is p -divisible. Then by part 1) of Proposition 2.10 together with the equation $\Sigma_{\mathcal{E}} = \frac{1}{p-1}vp - v(\vartheta - K)$, condition a) is equivalent to the equation

$$\frac{p}{p-1}vp - pv(\eta - K) = \frac{1}{p-1}vp - v(\eta - K).$$

Multiplying both sides with -1 and then adding $\frac{p}{p-1}vp$ to both sides, we find that this equation is equivalent to condition e).

Finally, if (K, v) has rank 1, then it follows as in the proof of Theorem 1.7 that condition d) is equivalent to condition g).

REFERENCES

- [1] Abhyankar, S.: *Two notes on formal power series*, Proc. Amer. Math. Soc. **7**, 903–905 (1956)
- [2] Atiyah, M.F. – MacDonald, I.G.: *Introduction to Commutative Algebra*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont. 1969
- [3] Blaszczyk, A. – Kuhlmann, F.-V.: *On maximal immediate extensions of valued fields*, Mathematische Nachrichten **290** (2017), 7–18
- [4] Coates, J. H. – Greenberg, R.: *Kummer theory for abelian varieties over local fields*, Invent. Math. **124** (1996), 129–174
- [5] Cutkosky, S.D.: *Introduction to Algebraic Geometry*, Graduate Studies in Mathematics **188**. American Mathematical Society, Providence, R.I., 2018
- [6] Cutkosky, S.D. – Kuhlmann, F.-V.: *Kähler differentials of extensions of valuation rings and deeply ramified fields*, arXiv:2306.04967
- [7] Silva de Souza, C.H. – Novacoski, J.H. – Spivakovskiy, M.: *Graded rings associated to valuations and direct limits*, Journal of Pure and Applied Algebra **227** (2023)
- [8] Eisenbud, D.: *Commutative Algebra with a view toward Algebraic Geometry*, Springer-Verlag, New York, 1995
- [9] Endler, O.: *Valuation theory*, Springer-Verlag, Berlin, 1972
- [10] Fesenko, I. B.: *On deeply ramified extensions*, J. London Math. Soc. (2) **57** (1998), 325–335
- [11] Gabber, O. – Ramero, L.: *Almost ring theory*, Lecture Notes in Mathematics **1800**, Springer-Verlag, Berlin, 2003

- [12] Jahnke, F. – Kartas, K.: *Beyond the Fontaine-Wintenberger theorem*, arXiv:2304.05881
- [13] Kaplansky, I.: *Maximal fields with valuations I*, Duke Math. Journ. **9** (1942), 303–321
- [14] Kato, K. – Kurokawa, N. – Saito, T.: *Number Theory 2: Introduction to Class Field Theory*, Translations of Mathematical Monographs **240**, 2011
- [15] Kuhlmann, F.-V.: *A classification of Artin–Schreier defect extensions and a characterization of defectless fields*, Illinois J. Math. **54** (2010), 397–448
- [16] Kuhlmann F.-V.: *The defect*, in: Commutative Algebra - Noetherian and non-Noetherian perspectives, Fontana, M., Kabbaj, S.-E., Olberding, B., Swanson, I. (Eds.), Springer-Verlag, New York, 2011
- [17] Kuhlmann, F.-V.: *Approximation of elements in henselizations*, manuscripta math. **136** (2011), 461–474
- [18] Kuhlmann, F.-V.: *Selected methods for the classification of cuts and their applications*, Proceedings of the ALANT 5 conference 2018, Banach Center Publications **121** (2020), 85–106
- [19] Kuhlmann, F.-V. – Kuhlmann, K.: *Arithmetic of cuts in ordered abelian groups and of ideals over valuation rings*, arXiv:2406.10545
- [20] Kuhlmann, F.-V. – Rzepka, A.: *The valuation theory of deeply ramified fields and its connection with defect extensions*, Transactions Amer. Math. Soc. **376** (2023), 2693–2738
- [21] Kuhlmann, F.-V. – Rzepka, A.: *The defect, II*, in preparation
- [22] Kuhlmann, F.-V. – Vlahu, I.: *The relative approximation degree*, Math. Z. **276** (2014), 203–235
- [23] Lang, S.: *Algebra*, revised 3rd. ed., Springer-Verlag, New York, 2002
- [24] Matsumura, H.: *Commutative Algebra, second Edition*, Mathematics Lecture Note Series, 56. Benjamin/Cummings Publishing Co., Inc., Reading, Mass., 1980
- [25] Novacoski, J. – Spivakovsky, M.: *Key polynomials and pseudo-convergent sequences*, J. Algebra **495** (2018), 199–219
- [26] Novacoski, J. – Spivakovsky, M.: *Kähler differentials, pure extensions and minimal key polynomials*, arXiv:2311.14322
- [27] Rotman, J.: *An Introduction to Homological Algebra*, Pure and Applied Mathematics, 85. Academic Press, Inc., New York-London, 1979
- [28] Tang, H.T.: *Gauss Lemma*, Proc. Amer. Math. Soc. **35**, 1972
- [29] Thatte, V.: *Ramification theory for Artin-Schreier extensions of valuation rings*, J. Algebra **456** (2016), 355–389
- [30] Thatte, V.: *Ramification theory for degree p extensions of valuation rings in mixed characteristic $(0, p)$* , J. Algebra **507** (2018), 225–248
- [31] Zariski, O. – Samuel, P.: *Commutative Algebra*, Vol. I, D. Van Nostrand, Princeton N.J., 1958
- [32] Zariski, O. – Samuel, P.: *Commutative Algebra*, Vol. II, New York–Heidelberg–Berlin, 1960

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA
Email address: cutkoskys@missouri.edu

INSTITUTE OF MATHEMATICS, UNIVERSITY OF SZCZECIN, UL. WIELKOPOLSKA 15, 70-451 SZCZECIN, POLAND
Email address: fvk@usz.edu.pl

INSTITUTE OF MATHEMATICS, UNIVERSITY OF SILESIA IN KATOWICE, BANKOWA 14, 40-007 KATOWICE, POLAND
Email address: anna.rzepka@us.edu.pl